

**ARG10020**  
anti-FGF basic antibody [F-474]Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [F-474] recognizes FGF basic
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Mouse
Clonality	Monoclonal
Clone	F-474
Isotype	IgG2a, kappa
Target Name	FGF basic
Antigen Species	Human
Immunogen	Purified recombinant human bFGF
Conjugation	Un-conjugated
Alternate Names	FGF-2; Fibroblast growth factor 2; bFGF; FGFB; Heparin-binding growth factor 2; BFGF; HBGF-2; Basic fibroblast growth factor

### Application Instructions

Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.
Calculated Mw	31 kDa

### Properties

Form	Liquid
Purification	Protein G affinity purified
Buffer	0.01M PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation


Database links	<a href="#">GeneID: 2247 Human</a>
----------------	------------------------------------

[Swiss-port # P09038 Human](#)

Gene Symbol	FGF2
Gene Full Name	fibroblast growth factor 2 (basic)
Background	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF. [provided by RefSeq, Jul 2008]
Function	Plays an important role in the regulation of cell survival, cell division, angiogenesis, cell differentiation and cell migration. Functions as potent mitogen in vitro. [UniProt]
Highlight	Related products: <a href="#">FGF basic antibodies</a> ; <a href="#">FGF basic ELISA Kits</a> ; <a href="#">FGF basic recombinant proteins</a> ; <a href="#">Anti-Mouse IgG secondary antibodies</a> ; Related news: <a href="#">The role of HDGF in tumor angiogenesis</a>
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Developmental Biology antibody; Neuroscience antibody; Signaling Transduction antibody

## Images

---



ARG10020 anti-FGF basic antibody [F-474] WB image

Western Blot: Recombinant human bFGF stained with anti-FGF basic antibody [F-474] (ARG10020)