

## ARG10088 anti-GCPII / PSMA antibody [YPSMA-1]

Package: 250 µg, 125 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [YPSMA-1] recognizes GCPII / PSMA
Tested Reactivity	Hu
Tested Application	ELISA, FACS, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	Little or no cross-reactivity to benign prostate hyperplasia (BPH) or to normal prostatic tissue.
Host	Mouse
Clonality	Monoclonal
Clone	YPSMA-1
Isotype	IgG2b, kappa
Target Name	GCPII / PSMA
Species	Human
Immunogen	Crude membrane protein preparation from pooled prostate malignant carcinoma from China.
Conjugation	Un-conjugated
Alternate Names	FOLH1; Folate Hydrolase 1; NAALAD1; GCPII; PSMA; PSM; Glutamate Carboxypeptidase 2; GCP2; FOLH; N-Acetylated-Alpha-Linked Acidic Dipeptidase I; Pteroylpoly-Gamma-Glutamate Carboxypeptidase; Folylpoly-Gamma-Glutamate Carboxypeptidase; Cell Growth-Inhibiting Gene 27 Protein; Membrane Glutamate Carboxypeptidase; Glutamate Carboxypeptidase II; Glutamate Carboxylase II; EC 3.4.17.21; NAALADase I; NAALAdase; FGCP; MGCP; Folate Hydrolase (Prostate-Specific Membrane Antigen) 1; N-Acetylated Alpha-Linked Acidic Dipeptidase 1; Prostate-Specific Membrane Antigen

### Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	FACS	Assay-dependent
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	Assay-dependent
	WB	Assay-dependent
Application Note	Immuno-histochemistry: This antibody was used on formalin-fixed paraffin embedded and frozen tissue sections.	
	Indirect ELISA: This antibody is reactive to PSMA expressing LNCap cells coated ELISA plate.	
	Western blot: The antibody, when used at 1µg/ml concentration, identified the 100KDa PSMA in LNCap cell extract. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

## Properties

Form	Liquid
Purification	Protein G affinity purified
Buffer	0.01M PBS (pH 7.0)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links	<a href="#">GeneID: 2346 Human</a> <a href="#">Swiss-port # Q04609 Human</a>
Gene Symbol	FOLH1
Gene Full Name	folate hydrolase (prostate-specific membrane antigen) 1
Background	This gene encodes a type II transmembrane glycoprotein belonging to the M28 peptidase family. The protein acts as a glutamate carboxypeptidase on different alternative substrates, including the nutrient folate and the neuropeptide N-acetyl-L-aspartyl-L-glutamate and is expressed in a number of tissues such as prostate, central and peripheral nervous system and kidney. A mutation in this gene may be associated with impaired intestinal absorption of dietary folates, resulting in low blood folate levels and consequent hyperhomocysteinemia. Expression of this protein in the brain may be involved in a number of pathological conditions associated with glutamate excitotoxicity. In the prostate the protein is up-regulated in cancerous cells and is used as an effective diagnostic and prognostic indicator of prostate cancer. This gene likely arose from a duplication event of a nearby chromosomal region. Alternative splicing gives rise to multiple transcript variants encoding several different isoforms.
Function	Has both folate hydrolase and N-acetylated-alpha-linked-acidic dipeptidase (NAALADase) activity. Has a preference for tri-alpha-glutamate peptides. In the intestine, required for the uptake of folate. In the brain, modulates excitatory neurotransmission through the hydrolysis of the neuropeptide, N-acetylaspartylglutamate (NAAG), thereby releasing glutamate. Involved in prostate tumor progression.
Research Area	Cancer antibody; Metabolism antibody; Signaling Transduction antibody
Calculated Mw	84 kDa
PTM	Glycoprotein, Phosphoprotein
Cellular Localization	Cell membrane, Cytoplasm, Membrane