

ARG10184 anti-beta Amyloid (1 - 42) antibody [CA9 10C11] (Biotin)

Package: 100 µl
Store at: -20°C

Summary

Product Description	Biotin-conjugated Mouse Monoclonal antibody [CA9 10C11] recognizes beta Amyloid (1 - 42)
Tested Reactivity	Hu, Ms, Rat, NHuPrm
Tested Application	ELISA
Specificity	Does not cross react with amyloid beta peptide 40 in dot blotting and ELISA. Cross-reactivity to amyloid beta peptide 43 is less than 1% in ELISA.
Host	Mouse
Clonality	Monoclonal
Clone	CA9 10C11
Isotype	IgG2b, kappa
Target Name	beta Amyloid (1 - 42)
Immunogen	KLH conjugated to a short peptide (MVGGVVIA) with amino acid sequence corresponding to the C-terminal of Aβ42
Conjugation	Biotin
Alternate Names	CVAP; AAA; AICD-50; PN2; 50; Beta-APP42; AID; Gamma-CTF; S-APP-alpha; 57; AD1; PN-II; Beta-APP40; 42; 40; APPI; Alzheimer disease amyloid protein; Amyloid beta A4 protein; PreA4; ABETA; Amyloid intracellular domain 50; CTFgamma; Amyloid intracellular domain 57; 59; AICD-59; S-APP-beta; APP; AICD-57; Amyloid intracellular domain 59; ABPP; Protease nexin-II; Cerebral vascular amyloid peptide

Application Instructions

Application table	Application	Dilution
	ELISA	1:10000

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Protein G affinity purified
Buffer	0.01M PBS (pH 7.0), 1% Gelatin and 0.1% Proclin-300
Preservative	0.1% Proclin-300
Stabilizer	1% Gelatin
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Keep the antibody in the dark and keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol	App
Gene Full Name	amyloid beta (A4) precursor protein
Background	<p>Amyloid beta peptide 42 (Aβ42) is best known for its role in the formation of senile plaques in the brain of patients with Alzheimer's disease. Aβ42 and Aβ40 are the two major amyloid peptides that are produced after cleavage of amyloid precursor protein by secretases. Aβ42 (42 amino acids) is very fibrillogenic. The beta pleated structure of Aβ42 constitutes the initial and key component of the insoluble amyloid fibril in senile plaque. It is widely accepted that Aβ42 contributes to the pathogenesis of Alzheimer's disease. One proposition is that the deposition of amyloid fibril onto the brain tissue results in Alzheimer's disease. Another is that the neurotoxicity of Aβ42 oligomer is the cause of the disease.</p>
Function	<p>Functions as a cell surface receptor and performs physiological functions on the surface of neurons relevant to neurite growth, neuronal adhesion and axonogenesis. Involved in cell mobility and transcription regulation through protein-protein interactions (By similarity). Can promote transcription activation through binding to APBB1-KAT5 and inhibit Notch signaling through interaction with Numb (By similarity). Couples to apoptosis-inducing pathways such as those mediated by G(O) and JIP. Inhibits G(o) alpha ATPase activity. Acts as a kinesin I membrane receptor, mediating the axonal transport of beta-secretase and presenilin 1 (By similarity). May be involved in copper homeostasis/oxidative stress through copper ion reduction. Can regulate neurite outgrowth through binding to components of the extracellular matrix such as heparin and collagen I and IV (By similarity). The splice isoforms that contain the BPTI domain possess protease inhibitor activity. Induces a AGER-dependent pathway that involves activation of p38 MAPK, resulting in internalization of amyloid-beta peptide and leading to mitochondrial dysfunction in cultured cortical neurons. Provides Cu(2+) ions for GPC1 which are required for release of nitric oxide (NO) and subsequent degradation of the heparan sulfate chains on GPC1 (By similarity).</p> <p>Beta-amyloid peptides are lipophilic metal chelators with metal-reducing activity. Binds transient metals such as copper, zinc and iron. Rat and mouse beta-amyloid peptides bind only weakly transient metals and have little reducing activity due to substitutions of transient metal chelating residues. Beta-APP42 may activate mononuclear phagocytes in the brain and elicits inflammatory responses. Promotes both tau aggregation and TPK II-mediated phosphorylation. Also bind GPC1 in lipid rafts (By similarity). Appicans elicit adhesion of neural cells to the extracellular matrix and may regulate neurite outgrowth in the brain.</p> <p>The gamma-CTF peptides as well as the caspase-cleaved peptides, including C31, are potent enhancers of neuronal apoptosis.</p> <p>N-APP binds TNFRSF21 triggering caspase activation and degeneration of both neuronal cell bodies (via caspase-3) and axons (via caspase-6). [UniProt]</p>
Highlight	<p>Related Antibody Duos and Panels: ARG30063 Beta amyloid peptide 42 ELISA Antibody Duo</p> <p>Related products: beta Amyloid antibodies; beta Amyloid ELISA Kits; beta Amyloid Duos / Panels; Anti-Mouse IgG secondary antibodies;</p>
Research Area	Neuroscience antibody
Calculated Mw	87 kDa. (79 - 120 kDa depending on glycosylation level)
PTM	<p>Proteolytically processed under normal cellular conditions. Cleavage either by alpha-secretase, beta-secretase or theta-secretase leads to generation and extracellular release of soluble APP peptides, S-APP-alpha and S-APP-beta, and the retention of corresponding membrane-anchored C-terminal fragments, C80, C83 and C99. Subsequent processing of C80 and C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway and is non-amyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59). Many other minor beta-amyloid peptides, beta-amyloid 1-X peptides, are found in cerebral spinal fluid (CSF) including the beta-amyloid X-15 peptides, produced from the cleavage by alpha-secretase and all terminating at Gln-686.</p> <p>Proteolytically cleaved by caspases during neuronal apoptosis. Cleavage at Asp-739 by either caspase-6, -8 or -9 results in the production of the neurotoxic C31 peptide and the increased production of beta-amyloid peptides.</p> <p>N- and O-glycosylated. O-glycosylation on Ser and Thr residues with core 1 or possibly core 8 glycans. Partial tyrosine glycosylation (Tyr-681) is found on some minor, short beta-amyloid peptides (beta-amyloid 1-15, 1-16, 1-17, 1-18, 1-19 and 1-20) but not found on beta-amyloid 38, beta-amyloid 40 nor on beta-amyloid 42. Modification on a tyrosine is unusual and is more prevalent in AD patients. Glycans had Neu5AcHex(Neu5Ac)HexNAc-O-Tyr, Neu5AcNeu5AcHex(Neu5Ac)HexNAc-O-Tyr and O-AcNeu5AcNeu5AcHex(Neu5Ac)HexNAc-O-Tyr structures, where O-Ac is O-acetylation of Neu5Ac. Neu5AcNeu5Ac is most likely Neu5Ac 2,8Neu5Ac linked. O-glycosylations in the vicinity of the cleavage</p>

sites may influence the proteolytic processing. Appicans are L-APP isoforms with O-linked chondroitin sulfate.

Phosphorylation in the C-terminal on tyrosine, threonine and serine residues is neuron-specific.

Phosphorylation can affect APP processing, neuronal differentiation and interaction with other proteins. Phosphorylated on Thr-743 in neuronal cells by Cdc5 kinase and Mapk10, in dividing cells by Cdc2 kinase in a cell-cycle dependent manner with maximal levels at the G2/M phase and, in vitro, by GSK-3-beta. The Thr-743 phosphorylated form causes a conformational change which reduces binding of Fe65 family members. Phosphorylation on Tyr-757 is required for SHC binding. Phosphorylated in the extracellular domain by casein kinases on both soluble and membrane-bound APP. This phosphorylation is inhibited by heparin.

Extracellular binding and reduction of copper, results in a corresponding oxidation of Cys-144 and Cys-158, and the formation of a disulfide bond. In vitro, the APP-Cu(+) complex in the presence of hydrogen peroxide results in an increased production of beta-amyloid-containing peptides.

Trophic-factor deprivation triggers the cleavage of surface APP by beta-secretase to release sAPP-beta which is further cleaved to release an N-terminal fragment of APP (N-APP).

Beta-amyloid peptides are degraded by IDE.