

## ARG10244 anti-Insulin antibody [C7C9]

Package: 100 µg, 50 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [C7C9] recognizes Insulin
Tested Reactivity	Hu, Bov, Pig
Tested Application	ELISA
Host	Mouse
Clonality	Monoclonal
Clone	C7C9
Isotype	IgG1
Target Name	Insulin
Immunogen	purified human insulin
Conjugation	Un-conjugated
Alternate Names	IDDM; IDDM2; IDDM1; ILPR; MODY10; Insulin; IRDN

### Application Instructions

Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.
Calculated Mw	12 kDa

### Properties

Form	Liquid
Purification	Protein A affinity purified.
Buffer	PBS (pH 7.4) and 0.1% Sodium azide
Preservative	0.1% Sodium azide
Concentration	1.0-2.0 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links	<a href="#">GeneID: 3630 Human</a> <a href="#">GeneID: 397415 Pig</a>
----------------	--------------------------------------------------------------------------

[Swiss-port # P01308 Human](#)

[Swiss-port # P01315 Pig](#)

<b>Gene Symbol</b>	INS
<b>Gene Full Name</b>	insulin
<b>Background</b>	After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2010]
<b>Function</b>	Insulin decreases blood glucose concentration. It increases cell permeability to monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in liver. [UniProt]
<b>Research Area</b>	Cell Biology and Cellular Response antibody; Metabolism antibody; Neuroscience antibody; Signaling Transduction antibody