

ARG10366 anti-H-FABP / Cardiac FABP antibody [9F3]

Package: 200 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [9F3] recognizes H-FABP / Cardiac FABP
Tested Reactivity	Hu
Tested Application	ELISA, Puri
Host	Mouse
Clonality	Monoclonal
Clone	9F3
Isotype	IgG1
Target Name	H-FABP / Cardiac FABP
Species	Human
Immunogen	human heart FABP
Conjugation	Un-conjugated
Alternate Names	FABP11; H-FABP; O-FABP; Heart-type fatty acid-binding protein; MDGI; Fatty acid-binding protein 3; Muscle fatty acid-binding protein; Mammary-derived growth inhibitor; Fatty acid-binding protein, heart; M-FABP

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	Puri	Assay-dependent
Application Note	Sandwich ELISA (Capture antibody - Detection antibody): ARG10366 - ARG10364 * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Protein A affinity purified.
Buffer	PBS (pH 7.4) and 0.1% Sodium azide
Preservative	0.1% Sodium azide
Concentration	1.0-2.0 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 2170 Human](#)

[Swiss-port # P05413 Human](#)

Gene Symbol

FABP3

Gene Full Name

fatty acid binding protein 3, muscle and heart

Background

The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is a candidate tumor suppressor gene for human breast cancer. [provided by RefSeq, Jul 2008]

Function

FABP are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters. [UniProt]


Research Area

Cell Biology and Cellular Response antibody; Developmental Biology antibody; Metabolism antibody

Calculated Mw

15 kDa

Images



ARG10366 anti-H-FABP / Cardiac FABP antibody [9F3] ELISA image

ELISA: Interaction of ARG10366 anti-H-FABP / Cardiac FABP antibody [9F3] with H-FABP, isolated from the Human heart for in direct ELISA.