

ARG10369 anti-Fibrinogen antibody [49D2]

Package: 200 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [49D2] recognizes Fibrinogen
Tested Reactivity	Hu
Tested Application	EIA, ELISA, WB
Host	Mouse
Clonality	Monoclonal
Clone	49D2
Isotype	IgG2a
Target Name	Fibrinogen
Antigen Species	Human
Immunogen	fibrinopeptide A conjugated with a carrier protein
Conjugation	Un-conjugated
Alternate Names	Fibrinogen alpha chain; Fib2

Application Instructions

Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.
Calculated Mw	95 kDa

Properties

Form	Liquid
Purification	Protein A affinity purified.
Buffer	PBS (pH 7.4) and 0.1% Sodium azide
Preservative	0.1% Sodium azide
Concentration	1.0-2.0 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 2243 Human
----------------	------------------------------------

Gene Symbol	FGA
Gene Full Name	fibrinogen alpha chain
Background	<p>The protein encoded by this gene is the alpha component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia, afibrinogenemia and renal amyloidosis. Alternative splicing results in two isoforms which vary in the carboxy-terminus. [provided by RefSeq, Jul 2008]</p>
Function	<p>Cleaved by the protease thrombin to yield monomers which, together with fibrinogen beta (FGB) and fibrinogen gamma (FGG), polymerize to form an insoluble fibrin matrix. Fibrin has a major function in hemostasis as one of the primary components of blood clots. In addition, functions during the early stages of wound repair to stabilize the lesion and guide cell migration during re-epithelialization. Was originally thought to be essential for platelet aggregation, based on in vitro studies using anticoagulated blood. However, subsequent studies have shown that it is not absolutely required for thrombus formation in vivo. Enhances expression of SELP in activated platelets via an ITGB3-dependent pathway. Maternal fibrinogen is essential for successful pregnancy. Fibrin deposition is also associated with infection, where it protects against IFNG-mediated hemorrhage. May also facilitate the immune response via both innate and T-cell mediated pathways. [UniProt]</p>
Research Area	Cell Biology and Cellular Response antibody; Metabolism antibody