

ARG10515 anti-HIF-1 alpha antibody [ESEE122]

Package: 50 µg, µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [ESEE122] recognizes HIF-1 alpha
Tested Reactivity	Hu, Ms, Rat, Bov
Tested Application	ELISA, FACS, ICC/IF, IHC-Fr, IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	ESEE122
Isotype	IgG1
Target Name	HIF-1 alpha
Species	Human
Immunogen	Synthetic peptide within 329 to 530 aa of Human HIF-1A protein (NP_001230013.1)
Conjugation	Un-conjugated
Alternate Names	Class E basic helix-loop-helix protein 78; Basic-helix-loop-helix-PAS protein MOP1; Hypoxia-inducible factor 1-alpha; PAS domain-containing protein 8; HIF1-alpha; HIF1-ALPHA; HIF1; MOP1; HIF-1-alpha; PASD8; HIF-1A; HIF-1alpha; Member of PAS protein 1; ARNT-interacting protein; bHLHe78

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent.
	FACS	1µg for 10 ⁶ cells.
	ICC/IF	5 - 12 µg/ml
	IHC-Fr	1:1000 - 1:8000
	IHC-P	1:1000 - 1:8000
	WB	1:100 - 1:1000
Application Note	IHC-P: Antigen Retrieval: By heat mediation. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Hypoxic cell pellets.	
Observed Size	~130 kDa	

Properties

Form	Liquid
Purification	Purified by affinity chromatography.


Buffer	PBS and 0.02% Sodium azide
Preservative	0.02% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	HIF1A
Gene Full Name	hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)
Background	The alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1), which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2011]
Function	Functions as a master transcriptional regulator of the adaptive response to hypoxia. Under hypoxic conditions, activates the transcription of over 40 genes, including erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, HILPDA, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. Plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Binds to core DNA sequence 5'-[AG]CGTG-3' within the hypoxia response element (HRE) of target gene promoters. Activation requires recruitment of transcriptional coactivators such as CREBBP and EP300. Activity is enhanced by interaction with both, NCOA1 or NCOA2. Interaction with redox regulatory protein APEX seems to activate CTAD and potentiates activation by NCOA1 and CREBBP. Involved in the axonal distribution and transport of mitochondria in neurons during hypoxia. [Uniprot]
Highlight	Related products: HIF1 alpha antibodies ; HIF1 alpha ELISA Kits ; Anti-Mouse IgG secondary antibodies ; Related news: Baking soda restores circadian clock in tumor cells Hypoxia-induced transcription, histone demethylases are involved
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Gene Regulation antibody; Metabolism antibody
Calculated Mw	93 kDa
PTM	In normoxia, is hydroxylated on Pro-402 and Pro-564 in the oxygen-dependent degradation domain (ODD) by EGLN1/PHD2 and EGLN2/PHD1. EGLN3/PHD3 has also been shown to hydroxylate Pro-564. The hydroxylated prolines promote interaction with VHL, initiating rapid ubiquitination and subsequent proteasomal degradation. Deubiquitinated by USP20. Under hypoxia, proline hydroxylation is impaired and ubiquitination is attenuated, resulting in stabilization. In normoxia, is hydroxylated on Asn-803 by HIF1AN, thus abrogating interaction with CREBBP and EP300 and preventing transcriptional activation. This hydroxylation is inhibited by the Cu/Zn-chelator, Clioquinol. S-nitrosylation of Cys-800 may be responsible for increased recruitment of p300 coactivator necessary for transcriptional activity of HIF-1 complex. Requires phosphorylation for DNA-binding. Phosphorylation at Ser-247 by CSNK1D/CK1 represses kinase activity and impairs ARNT binding. Phosphorylation by GSK3-beta and PLK3 promote degradation by the proteasome. Sumoylated; with SUMO1 under hypoxia. Sumoylation is enhanced through interaction with RWDD3. Both sumoylation and desumoylation seem to be involved in the regulation of its stability during hypoxia. Sumoylation can promote either its stabilization or its VHL-dependent degradation by promoting hydroxyproline-independent HIF1A-VHL complex binding, thus leading to HIF1A ubiquitination and proteasomal degradation. Desumoylation by SENP1 increases its stability and transcriptional activity. There is a discord between various publications on the effect of sumoylation and desumoylation on its stability and transcriptional activity. Acetylation of Lys-532 by ARD1 increases interaction with VHL and stimulates subsequent proteasomal degradation (PubMed:12464182). Deacetylation of Lys-709 by SIRT2 increases its interaction with and hydroxylation by EGLN1 thereby inactivating HIF1A activity by inducing its proteasomal degradation (PubMed:24681946).


Polyubiquitinated; in normoxia, following hydroxylation and interaction with VHL. Lys-532 appears to be the principal site of ubiquitination. Clioquinol, the Cu/Zn-chelator, inhibits ubiquitination through preventing hydroxylation at Asn-803. Ubiquitinated by a CUL2-based E3 ligase. The iron and 2-oxoglutarate dependent 3-hydroxylation of asparagine is (S) stereospecific within HIF CTAD domains.

Images




ARG10515 anti-HIF-1 alpha antibody [ESEE122] IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse muscle tissue stained with ARG10515 anti-HIF-1 alpha antibody [ESEE122] at 1:90 dilution for 2 hours at 37°C.



ARG10515 anti-HIF-1 alpha [ESEE122] antibody WB image

Western blot: Whole cell extracts from HeLa (normoxia) or HeLa (0.1% hypoxia for 4 hours) stained with ARG10515 anti-HIF-1 alpha [ESEE122] antibody at 4 µg/ml dilution. (Clone WB data from Am J Pathol. 2000 Aug; 157(2): 411–421; PMID: 10934146)



ARG10515 anti-HIF-1 alpha antibody [ESEE122] ICC/IF image

Immunofluorescence: NRK-49F cells untreated or treated with CoCl₂ (400 µM) for 8 hours. Cells were fixed with 4% PFA for 15 min at room temperature and permeabilized by 0.5% Triton X-100 for 10 min at room temperature. Blocking: CAS-Blocking buffer for 1 hour. Cells were stained with ARG10515 anti-HIF-1 alpha antibody [ESEE122] (red) at 1:200 dilution, overnight at 4°C, in filtered PBS. DAPI (blue) for nuclear staining.