

## ARG10537 Goat anti-Mouse IgG (Fab) antibody (TRITC)

Package: 250 µl

Store at: -20°C

### Summary

Product Description	TRITC-conjugated Goat Polyclonal antibody recognizes Mouse IgG (Fab)
Tested Reactivity	Ms, Rat, Gpig
Tested Application	ELISA, ICC/IF, IHC-Fr
Specificity	TRITC-conjugated IgG fraction of polyclonal Goat antiserum to Mouse Fab of IgG .
Host	Goat
Clonality	Polyclonal
Target Name	Mouse IgG Fab
Target Ig	IgG
Conjugation	TRITC

### Application Instructions

Cross Reactivity Note	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has been tested in double radial immunodiffusion with the following results: Bovine (-), Duck (-), Human (-), Rat (+), Cat (-), Goat (±), Horse (-), Sheep (-), Chicken (-), Guinea Pig (+), Monkey (-), Swine (-), Dog (-), Hamster (-), Rabbit (-), Turkey (-).								
Species Does Not React With	Hu, Bov, Cat, Chk, Dog, Hm, Hrs, Mk, Pig, Rb, Sheep								
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>1:20 - 1:80</td></tr><tr><td>ICC/IF</td><td>1:20 - 1:80</td></tr><tr><td>IHC-Fr</td><td>1:20 - 1:80</td></tr></tbody></table>	Application	Dilution	ELISA	1:20 - 1:80	ICC/IF	1:20 - 1:80	IHC-Fr	1:20 - 1:80
Application	Dilution								
ELISA	1:20 - 1:80								
ICC/IF	1:20 - 1:80								
IHC-Fr	1:20 - 1:80								
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.								

### Properties

Form	Liquid
Buffer	PBS (pH 7.2)
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Keep the antibody in the dark and keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

