

ARG10581 anti-AMACR / p504S antibody [Q17-L]

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Monoclonal antibody [Q17-L] recognizes AMACR / p504S
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Monoclonal
Clone	Q17-L
Isotype	IgG
Target Name	AMACR / p504S
Species	Human
Immunogen	Synthetic peptide around the N-terminus of Human P504S
Conjugation	Un-conjugated
Alternate Names	CBAS4; RACE; RM; EC 5.1.99.4; Alpha-methylacyl-CoA racemase; AMACRD; 2-methylacyl-CoA racemase

Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>IHC-P</td><td>1:100 - 1:200</td></tr></tbody></table>	Application	Dilution	IHC-P	1:100 - 1:200
Application	Dilution				
IHC-P	1:100 - 1:200				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				

Properties


Form	Liquid
Buffer	20 mM Tris-HCl (pH 8.0), 0.05% Sodium azide and 20 mg/ml BSA
Preservative	0.05% Sodium azide
Stabilizer	20 mg/ml BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 23600 Human
----------------	-------------------------------------


Gene Symbol	AMACR
Gene Full Name	alpha-methylacyl-CoA racemase
Background	This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene. [provided by RefSeq, Mar 2011]
Function	Racemization of 2-methyl-branched fatty acid CoA esters. Responsible for the conversion of pristanoyl-CoA and C27-bile acyl-CoAs to their (S)-stereoisomers. [UniProt]
Calculated Mw	42 kDa

Images



ARG10581 anti-AMACR / p504S antibody [Q17-L] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human papillary renal cell carcinoma (4 μ m sections) stained with ARG10581 anti-AMACR / p504S antibody [Q17-L].



ARG10581 anti-AMACR / p504S antibody [Q17-L] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human prostate adenocarcinoma (4 μ m sections) stained with ARG10581 anti-AMACR / p504S antibody [Q17-L].