

ARG10658 anti-VEGF antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes VEGF
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	VEGF
Antigen Species	Human
Immunogen	Synthetic peptide around the N-terminus of Human VEGF (VDIFQEYPDEIEYIFK). (100% homologous in Human, Mouse and Rat).
Conjugation	Un-conjugated
Alternate Names	MVCD1; Vascular permeability factor; VEGF-A; VPF; VEGF; Vascular endothelial growth factor A

Application Instructions

Application table	Application	Dilution
	IHC-P	0.5 - 1 µg/ml
	WB	0.5 - 1 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	27 kDa	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.025% Sodium azide and 2.5% BSA.
Preservative	0.025% Sodium azide
Stabilizer	2.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation


Gene Symbol	VEGFA
Gene Full Name	vascular endothelial growth factor A
Background	This gene is a member of the PDGF/VEGF growth factor family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects, including mediating increased vascular permeability, inducing angiogenesis, vasculogenesis and endothelial cell growth, promoting cell migration, and inhibiting apoptosis. Elevated levels of this protein is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy. Alternatively spliced transcript variants, encoding either freely secreted or cell-associated isoforms, have been characterized. There is also evidence for the use of non-AUG (CUG) translation initiation sites upstream of, and in-frame with the first AUG, leading to additional isoforms. [provided by RefSeq, Jul 2008]
Function	Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth. [UniProt]
Highlight	Related products: VEGF antibodies ; VEGF ELISA Kits ; VEGF Duos / Panels ; VEGF recombinant proteins ; Anti-Rabbit IgG secondary antibodies ; Related news: The role of HDGF in tumor angiogenesis

Images



ARG10658 anti-VEGF antibody IHC-P image

Immunohistochemistry: Human lung cancer tissue stained with ARG10658 anti-VEGF antibody.



ARG10658 anti-VEGF antibody WB image

Western blot: 1) 10 ng, 2) 5 ng, and 3) 2.5 ng of recombinant Human protein stained with ARG10658 anti-VEGF antibody.