

## ARG10699 anti-Aurora (pan) antibody [4A7]

Package: 50 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [4A7] recognizes Aurora (pan)
Tested Reactivity	Hu, Ms, Rat, Cow, Hrs, Pig
Predict Reactivity	Chk
Tested Application	ICC/IF, IHC-Fr, WB
Host	Mouse
Clonality	Monoclonal
Clone	4A7
Isotype	IgG1
Target Name	Aurora (pan)
Antigen Species	Human
Immunogen	Full length recombinant Human Aurora C protein expressed in and purified from E. coli.
Conjugation	Un-conjugated
Alternate Names	ARK-1; AIK; BTAK; Serine/threonine-protein kinase 6; Breast tumor-amplified kinase; Serine/threonine-protein kinase aurora-A; STK15; Serine/threonine-protein kinase 15; AURORA2; Aurora-related kinase 1; hARK1; AURA; STK6; STK7; Aurora kinase A; EC 2.7.11.1; Aurora/IPL1-related kinase 1; Aurora 2; ARK1; PPP1R47

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:1000 - 1:2000
	IHC-Fr	1:1000 - 1:2000
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	46 kDa	

### Properties


Form	Liquid
Purification	Affinity purification.
Buffer	PBS and 50% Glycerol.
Stabilizer	50% Glycerol
Concentration	1 mg/ml

<b>Storage instruction</b>	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
<b>Note</b>	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


<b>Gene Symbol</b>	AURKA
<b>Gene Full Name</b>	aurora kinase A
<b>Background</b>	The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008]
<b>Function</b>	Mitotic serine/threonine kinases that contributes to the regulation of cell cycle progression. Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis. Required for initial activation of CDK1 at centrosomes. Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2. Regulates KIF2A tubulin depolymerase activity. Required for normal axon formation. Plays a role in microtubule remodeling during neurite extension. Important for microtubule formation and/or stabilization. Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint-response pathways critical for oncogenic transformation of cells, by phosphorylating and stabilizing p53/TP53. Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity. Necessary for proper cilia disassembly prior to mitosis. [UniProt]

## Images



ARG10699 anti-Aurora (pan) antibody [4A7] ICC/IF image

Immunocytochemistry: HeLa cell cultures were stained with ARG10699 anti-Aurora (pan) antibody [4A7] (green). In this image, strong staining in spindle poles was seen in cells at anaphase. This antibody also stains the midbodies between the two daughter cells. Cells were co-stained with a chicken polyclonal antibody to Vimentin (red). Blue is a DNA stain.



ARG10699 anti-Aurora (pan) antibody [4A7] WB image

Western blot: HeLa cells treated with 100 ng/ml nocodazole for 18 hours was stained with ARG10699 anti-Aurora (pan) antibody [4A7] (Left). Nocodazole is a microtubule depolymerizing agent which induces cells to halt at the G2/M phase and also induces Aurora kinase expression. ARG10699 anti-Aurora (pan) antibody [4A7] binds strongly to bands at about 46, 38, and 35 kDa, corresponding to Aurora A, B, and C (Right).