

Product datasheet

info@arigobio.com

ARG10798 anti-PDE3B antibody

Package: 50 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes PDE3B

Tested Reactivity Hu, Ms, Rat, Chk, Mk

Tested Application Confocal, Dot, ELISA, ICC/IF, IHC-P, IP, WB

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name PDE3B
Species Human

Immunogen Synthetic peptide taken within aa. 1060-1110 from Human PDE3B.

Conjugation Un-conjugated

Alternate Names EC 3.1.4.17; HcGIP1; cGIPDE1; CGIPDE1; CGIPDE B; Cyclic GMP-inhibited phosphodiesterase B; cGMP-

inhibited 3',5'-cyclic phosphodiesterase B; CGIP1

Application Instructions

Application table	Application	Dilution
	Confocal	1:500
	Dot	1:10000
	ELISA	1:10000
	ICC/IF	1:500
	IHC-P	1:500
	IP	1:200
	WB	1:500
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Affinity purified.

Buffer Tris-Glycine Buffer (pH 7.4 - 7.8), Hepes, 0.02% Sodium azide, 30% Glycerol and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 30% Glycerol and 0.5% BSA

www.arigobio.com arigo.nuts about antibodies 1/2

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol Gene Full Name PDE3B

phosphodiesterase 3B, cGMP-inhibited


Function Cyclic nucleotide phosphodiesterase with a dual-specificity for the second messengers cAMP and cGMP,

which are key regulators of many important physiological processes. May play a role in fat metabolism. Regulates cAMP binding of RAPGEF3. Through simultaneous binding to RAPGEF3 and PIK3R6 assembles a signaling complex in which the PI3K gamma complex is activated by RAPGEF3 and which is involved in

angiogenesis. [UniProt]

Calculated Mw 124 kDa

Images



ARG10798 anti-PDE3B antibody WB image

Western blot: Recombinant PDE3B protein stained with ARG10798 anti-PDE3B antibody at 1:500 dilution.