

ARG10816 anti-MERTK phospho (Tyr749 / Tyr753 / Tyr754) antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes MERTK phospho (Tyr749 / Tyr753 / Tyr754)
Tested Reactivity	Hu, Ms, Rat, Mk
Tested Application	Confocal, Dot, ELISA, ICC/IF, IHC, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MERTK
Antigen Species	Human
Immunogen	Phosphospecific peptide corresponding to aa. 746-757 of Human MERTK. (CKKI-Y(p)-SGD-Y(p)-Y(p)-RQG)
Conjugation	Un-conjugated
Alternate Names	c-mer; Receptor tyrosine kinase MerTK; c-Eyk; Tyro12; MER; RP38; EC 2.7.10.1; Proto-oncogene c-Mer; Tyrosine-protein kinase Mer

Application Instructions

Application table	Application	Dilution
	Confocal	1:250
	Dot	1:10000
	ELISA	1:10000
	ICC/IF	1:250
	IHC	1:250
	WB	1:750
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	110 kDa	
Observed Size	~ 175 kDa	

Properties


Form	Liquid
Purification	Affinity purified.
Buffer	Tris-Glycine Buffer (pH 7.4 - 7.8), Hepes, 0.02% Sodium azide, 30% Glycerol and 0.5% BSA.

Preservative	0.02% Sodium azide
Stabilizer	30% Glycerol and 0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	MERTK
Gene Full Name	MER proto-oncogene, tyrosine kinase
Background	This gene is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. Mutations in this gene have been associated with disruption of the retinal pigment epithelium (RPE) phagocytosis pathway and onset of autosomal recessive retinitis pigmentosa (RP). [provided by RefSeq, Jul 2008]
Function	Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to several ligands including LGALS3, TUB, TULP1 or GAS6. Regulates many physiological processes including cell survival, migration, differentiation, and phagocytosis of apoptotic cells (efferocytosis). Ligand binding at the cell surface induces autophosphorylation of MERTK on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with GRB2 or PLCG2 and induces phosphorylation of MAPK1, MAPK2, FAK/PTK2 or RAC1. MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells, platelet aggregation, cytoskeleton reorganization and engulfment. Functions in the retinal pigment epithelium (RPE) as a regulator of rod outer segments fragments phagocytosis. Plays also an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3. [UniProt]

Images



ARG10816 anti-MERTK phospho (Tyr749 / Tyr753 / Tyr754) antibody WB image

Western blot: 1) Phosphorylated, and 2,3) non-phosphorylated MERTK was immunoprecipitated with ARG10784 anti-MERTK antibody and stained with ARG10816 anti-MERTK phospho (Tyr749 / Tyr753 / Tyr754) antibody. MW of MERTK is 174 kDa. Lower broad band is IgG from immunoprecipitation.