

## ARG11030 anti-Aryl Hydrocarbon Receptor antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes Aryl Hydrocarbon Receptor
Tested Reactivity	Ms
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Aryl Hydrocarbon Receptor
Species	Human
Immunogen	Synthetic peptide around the C-terminus of Mouse Aryl Hydrocarbon Receptor. (EAALKHEQIGHAQDV)
Conjugation	Un-conjugated
Alternate Names	Aryl hydrocarbon receptor; AhR; Ah receptor; bHLHe76; Class E basic helix-loop-helix protein 76

### Application Instructions

Application table	Application	Dilution
	IHC-P	1 - 5 µg/ml
	WB	0.5 - 1 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 95 kDa	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.025% Sodium azide and 2.5% BSA.
Preservative	0.025% Sodium azide
Stabilizer	2.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	AHR
Gene Full Name	aryl hydrocarbon receptor
Background	The protein encoded by this gene is a ligand-activated helix-loop-helix transcription factor involved in the regulation of biological responses to planar aromatic hydrocarbons. This receptor has been shown to regulate xenobiotic-metabolizing enzymes such as cytochrome P450. Before ligand binding, the encoded protein is sequestered in the cytoplasm; upon ligand binding, this protein moves to the nucleus and stimulates transcription of target genes. [provided by RefSeq, Sep 2015]
Function	Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons. Involved in cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues. Regulates the circadian clock by inhibiting the basal and circadian expression of the core circadian component PER1. Inhibits PER1 by repressing the CLOCK-ARNTL/BMAL1 heterodimer mediated transcriptional activation of PER1. [UniProt]
Calculated Mw	96 kDa

Images




ARG11030 anti-Aryl Hydrocarbon Receptor antibody IHC-P

Immunohistochemistry: Paraffin-embedded Mouse heart tissue stained with ARG11030 anti-Aryl Hydrocarbon Receptor antibody.



ARG11030 anti-Aryl Hydrocarbon Receptor antibody WB image

Western blot: 1) Mouse brain, 2) Mouse heart and 3) Mouse liver lysates stained with ARG11030 anti-Aryl Hydrocarbon Receptor antibody.



#### ARG11030 anti-Aryl Hydrocarbon Receptor antibody IHC-P

Immunohistochemistry: Paraffin-embedded Mouse brain tissue stained with ARG11030 anti-Aryl Hydrocarbon Receptor antibody.