

## ARG11127 anti-MARCKS antibody [5F9]

Package: 50 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [5F9] recognizes MARCKS
Tested Reactivity	Hu, Mk
Species Does Not React With	Ms, Rat
Tested Application	ICC/IF, IHC-Fr, WB
Host	Mouse
Clonality	Monoclonal
Clone	5F9
Isotype	IgG1
Target Name	MARCKS
Species	Human
Immunogen	Full-length Human MARCKS.
Conjugation	Un-conjugated
Alternate Names	MACS; 80K-L; Myristoylated alanine-rich C-kinase substrate; PKCSL; Protein kinase C substrate, 80 kDa protein, light chain; 80K-L protein; MARCKS; PRKCSL

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:2000
	IHC-Fr	1:2000
	WB	1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified
Buffer	PBS, 5 mM Sodium azide and 50% Glycerol.
Preservative	5 mM Sodium azide
Stabilizer	50% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw

cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


#### Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	MARCKS
Gene Full Name	myristoylated alanine-rich protein kinase C substrate
Background	The protein encoded by this gene is a substrate for protein kinase C. It is localized to the plasma membrane and is an actin filament crosslinking protein. Phosphorylation by protein kinase C or binding to calcium-calmodulin inhibits its association with actin and with the plasma membrane, leading to its presence in the cytoplasm. The protein is thought to be involved in cell motility, phagocytosis, membrane trafficking and mitogenesis. [provided by RefSeq, Jul 2008]
Function	MARCKS is the most prominent cellular substrate for protein kinase C. This protein binds calmodulin, actin, and synapsin. MARCKS is a filamentous (F) actin cross-linking protein. [UniProt]
Calculated Mw	32 kDa
PTM	Phosphorylation by PKC displaces MARCKS from the membrane. It also inhibits the F-actin cross-linking activity. [UniProt]
Cellular Localization	Cytoplasm, cytoskeleton. Membrane; Lipid-anchor. [UniProt]

## Images



ARG11127 anti-MARCKS antibody [5F9] ICC/IF image

Immunofluorescence: HeLa cells stained with ARG11127 anti-MARCKS antibody [5F9] (red) at 1:1000 dilution, and co-stained with anti-Vimentin antibody (green) at 1:10000 dilution. Hoechst (blue) for nuclear staining.



ARG11127 anti-MARCKS antibody [5F9] WB image

Western blot: NIH/3T3, HEK293, HeLa, SH-SY5Y, C6 and COS-1 cell lysates stained with ARG11127 anti-MARCKS antibody [5F9].

The antibody does not recognize the rodent form of the MARCKS protein.