

## ARG20920 anti-CD38 antibody [NIMR-5] (PE)

Package: 50 µg  
Store at: 4°C

### Summary

Product Description	PE-conjugated Rat Monoclonal antibody [NIMR-5] recognizes CD38
Tested Reactivity	Ms
Tested Application	BL, Cell-Act , FACS, IHC-Fr
Specificity	Mouse CD38.
Host	Rat
Clonality	Monoclonal
Clone	NIMR-5
Isotype	IgG2a, kappa
Target Name	CD38
Species	Mouse
Immunogen	BCL1 plasma membrane glycoproteins
Conjugation	PE
Alternate Names	cADPr hydrolase 1; ADPRC 1; EC 3.2.2.6; 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase; Cyclic ADP-ribose hydrolase 1; ADPRC1; EC 2.4.99.20; ADP-ribosyl cyclase 1; 2'-phospho-cyclic-ADP-ribose transferase; CD antigen CD38; T10; 2'-phospho-ADP-ribosyl cyclase; ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1

### Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Cell-Act	Assay-dependent
	FACS	< 0.2 µg/10 <sup>6</sup> cells
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Buffer	PBS, 0.1% Sodium azide and Sucrose.
Preservative	0.1% Sodium azide
Stabilizer	Sucrose
Concentration	0.1 mg/ml

Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links	<a href="#">GeneID: 12494 Mouse</a> <a href="#">Swiss-port # P56528 Mouse</a>
Gene Symbol	CD38
Gene Full Name	CD38 antigen
Background	The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]
Function	Synthesizes the second messengers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system. [UniProt]
Research Area	Cancer antibody; Developmental Biology antibody; Immune System antibody; Metabolism antibody; Pro-B Cell Marker antibody; Pre-B Cell Marker antibody
Calculated Mw	34 kDa

## Images



ARG20920 anti-CD38 antibody [NIMR-5] (PE) FACS image

Flow Cytometry: BALB/c Mouse splenocytes stained with ARG20920 anti-CD38 antibody [NIMR-5] (PE) and [ARG65532](#) anti-CD45R / B220 antibody [RA3-6B2] (FITC).