

ARG20944
anti-CD71 / Transferrin Receptor antibody [RI7217] (Biotin)Package: 100 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rat Monoclonal antibody [RI7217] recognizes CD71 / Transferrin Receptor
Tested Reactivity	Ms
Tested Application	BL, FACS, ICC/IF, WB
Specificity	Mouse CD71. The clone RI7217 inhibits cell proliferation in vitro.
Host	Rat
Clonality	Monoclonal
Clone	RI7217
Isotype	IgG2a, kappa
Target Name	CD71 / Transferrin Receptor
Antigen Species	Mouse
Immunogen	DMSO induced Friend erythroleukemia 745.6
Conjugation	Biotin
Alternate Names	TFR1; CD antigen CD71; CD71; T9; p90; TR; Trfr; Transferrin receptor protein 1; TRFR; sTfR; TfR1; TfR; TFR

Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	FACS	< 1 µg/10 ⁶ cells
	ICC/IF	Assay-dependent
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	85 kDa	

Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 22042 Mouse](#)

[Swiss-port # Q62351 Mouse](#)

Gene Symbol

TFRC

Gene Full Name

transferrin receptor


Background

This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeq, Sep 2015]

Function

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. [UniProt]

Images



ARG20944 anti-CD71 / Transferrin Receptor antibody [RI7217] (Biotin) FACS image

Flow Cytometry: Mouse pre-B cell line 18-81 was stained with ARG20944 anti-CD71 / Transferrin Receptor antibody [RI7217] (Biotin) followed by Streptavidin (FITC).