

ARG20985 anti-Ly6C antibody [HK1.4]

Package: 100 µg
Store at: -20°C

Summary

| | |
|---------------------|---|
| Product Description | Rat Monoclonal antibody [HK1.4] recognizes Ly6C |
| Tested Reactivity | Ms |
| Tested Application | Cell-Act , FACS, IHC-Fr |
| Specificity | Mouse Ly-6C. |
| Host | Rat |
| Clonality | Monoclonal |
| Clone | HK1.4 |
| Isotype | IgG2c, kappa |
| Target Name | Ly6C |
| Species | Mouse |
| Immunogen | L3 cloned CTL cells |
| Conjugation | Un-conjugated |
| Alternate Names | AA959465; Ly-6C1; AA682074; Ly-6C; Lymphocyte antigen 6C1; Ly6c |

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|------------------------------|
| | Cell-Act | Assay-dependent |
| | FACS | < 1 µg/10 ⁶ cells |
| | IHC-Fr | Assay-dependent |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

| | |
|---------------------|---|
| Form | Liquid |
| Buffer | BBS (pH 8.2) |
| Concentration | 0.5 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Bioinformation

| | |
|----------------|--|
| Database links | GeneID: 17067 Mouse Swiss-port # POCW02 Mouse |
| Gene Symbol | Ly6c1 |
| Gene Full Name | lymphocyte antigen 6 complex, locus C1 |
| Highlight | Related products: Ly6 antibodies ; Ly6 ELISA Kits ; Ly6 Duos / Panels ; Anti-Rat IgG secondary antibodies ; Related news: New antibody panels and duos for Tumor immune microenvironment Exploring Antiviral Immune Response |
| Research Area | Mouse Inflammatory Cell Marker antibody; Neutrophil Marker antibody; Mouse MDSC Marker antibody; Myeloid-derived suppressor cell antibody |
| Calculated Mw | 14 kDa |