

ARG21004 anti-CD152 / CTLA4 antibody [1B8] (PE)

Package: 50 µg
Store at: 4°C

Summary

Product Description	PE-conjugated Hamster Monoclonal antibody [1B8] recognizes CD152 / CTLA4
Tested Reactivity	Ms
Tested Application	FACS, FLISA
Specificity	Mouse CD152.
Host	Hamster
Clonality	Monoclonal
Clone	1B8
Isotype	IgG1
Target Name	CD152 / CTLA4
Species	Mouse
Immunogen	Extracellular portion of murine CTLA-4 fused to a murine IgG2a
Conjugation	PE
Alternate Names	GRD4; CTLA-4; CELIAC3; CD; Cytotoxic T-lymphocyte-associated antigen 4; CD152; GSE; CD antigen CD152; Cytotoxic T-lymphocyte protein 4; ALPS5; IDDM12

Application Instructions

Application table	Application	Dilution
	FACS	< 0.3 µg/10 ⁶ cells
	FLISA	
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Buffer	PBS, 0.1% Sodium azide and Sucrose.
Preservative	0.1% Sodium azide
Stabilizer	Sucrose
Concentration	0.1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 12477 Mouse Swiss-port # P09793 Mouse
Gene Symbol	CTLA4
Gene Full Name	cytotoxic T-lymphocyte-associated protein 4
Background	This gene is a member of the immunoglobulin superfamily and encodes a protein which transmits an inhibitory signal to T cells. The protein contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. Mutations in this gene have been associated with insulin-dependent diabetes mellitus, Graves disease, Hashimoto thyroiditis, celiac disease, systemic lupus erythematosus, thyroid-associated orbitopathy, and other autoimmune diseases. [provided by RefSeq, Jul 2008]
Function	Inhibitory receptor acting as a major negative regulator of T-cell responses. The affinity of CTLA4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28. [UniProt]
Calculated Mw	25 kDa
PTM	N-glycosylation is important for dimerization. Phosphorylation at Tyr-201 prevents binding to the AP-2 adapter complex, blocks endocytosis, and leads to retention of CTLA4 on the cell surface.

Images



ARG21004 anti-CD152 / CTLA4 antibody [1B8] (PE) FLISA image

FLISA: The plate was coated with Mouse CD152 / CTLA4 and purified/unlabeled Rat IgG1k. CTLA4 and purified immunoglobulin were detected with serially diluted ARG21004 anti-CD152 / CTLA4 antibody [1B8] (PE).