

## ARG21037 anti-CD81 antibody [2F7] (FITC)

Package: 100 µg  
Store at: 4°C

### Summary

Product Description	FITC-conjugated Hamster Monoclonal antibody [2F7] recognizes CD81
Tested Reactivity	Ms
Tested Application	BL, FACS, ICC/IF, IHC
Specificity	Mouse CD81. The clone 2F7 can block thymocyte interaction with CD81 in vitro.
Host	Hamster
Clonality	Monoclonal
Clone	2F7
Isotype	IgG3
Target Name	CD81
Antigen Species	Mouse
Immunogen	Mouse epithelial cell line PAM212
Conjugation	FITC
Alternate Names	CD antigen CD81; TAPA1; Tspan-28; S5.7; CD81 antigen; Target of the antiproliferative antibody 1; Tetraspanin-28; 26 kDa cell surface protein TAPA-1; CVID6; TSPAN28

### Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	FACS	< 1 µg/10 <sup>6</sup> cells
	ICC/IF	Assay-dependent
	IHC	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	26 kDa	

### Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links

[GeneID: 12520 Mouse](#)

[Swiss-port # P35762 Mouse](#)

Gene Symbol

CD81

Gene Full Name

CD81 antigen

Background


The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. This protein appears to promote muscle cell fusion and support myotube maintenance. Also it may be involved in signal transduction. This gene is localized in the tumor-suppressor gene region and thus it is a candidate gene for malignancies. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2014]

Function

May play an important role in the regulation of lymphoma cell growth. Interacts with a 16-kDa Leu-13 protein to form a complex possibly involved in signal transduction. May act as the viral receptor for HCV. [UniProt]

## Images

---



ARG21037 anti-CD81 antibody [2F7] (FITC) FACS image

Flow Cytometry: Chinese hamster ovary cell line CHO-K1 (negative control) and BALB/cAnN Mouse B lymphocyte cell line A20 stained with ARG21037 anti-CD81 antibody [2F7] (FITC).