

ARG21065 anti-MHC Class II I E kappa antibody [14-4-4S]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [14-4-4S] recognizes MHC Class II I E kappa
Tested Reactivity	Ms, Rat
Tested Application	BL, Cell-Act , FACS, ICC/IF, IHC-Fr, IP
Specificity	Mouse I-Ek/Rat RT1D. The clone 14-4-4S reacts with the I-Ek class II alloantigen on cells from mice of the H-2d, H-2p, and H-2r haplotypes. Cells from mice of the H-2b, H-2f, H-2q, and H-2s haplotypes do not express I-E antigen. The antibody has been reported to cross-react with the rat MHC class II alloantigen RT1D.
Host	Mouse
Clonality	Monoclonal
Clone	14-4-4S
Isotype	IgG2a, kappa
Target Name	MHC Class II I E kappa
Antigen Species	Mouse
Immunogen	C3H mouse skin graft and splenocytes
Conjugation	Un-conjugated
Alternate Names	A1323765; H-2Ea; MHC-H2-Ea; H2-Ea; I-Ealpha; H-2 class II histocompatibility antigen, E-U alpha chain; Ia3; E-alpha-f; Ia-3

Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Cell-Act	Assay-dependent
	FACS	Assay-dependent
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	29 kDa	

Properties

Form	Liquid
------	--------

Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 100504404 Mouse Swiss-port # P14439 Mouse
Gene Symbol	H2-Ea-ps
Gene Full Name	histocompatibility 2, class II antigen E alpha, pseudogene
Background	This locus belongs to the class II major histocompatibility complex (MHC) family of genes, which encode immune response (Ia) antigens that function in the T-cell-dependent immune response. This family member has multiple haplotypes, some of which result in the production of an E-alpha subunit that combines with an E-beta subunit to form a functional E complex at the cell surface. Other haplotypes, including that of the reference genome allele, contain mutations and they thus represent polymorphic pseudogenes that do not produce functional products. These mutations include frameshifting indels, nonsense mutations, and deletions of larger regions. The reference genome haplotype contains a deletion at the 5' end of the gene, including the core promoter region and the transcription start site, and therefore no transcripts result from this haplotype. [provided by RefSeq, Aug 2011]