

## ARG21075 anti-MHC Class I H2 Db antibody [28-14-8] (low endotoxin)

Package: 100 μg Store at: -20°C

Summary	
Product Description	Azide free and low endotoxin Mouse Monoclonal antibody [28-14-8] recognizes MHC Class I H2 Db
Tested Reactivity	Ms
Tested Application	BL, FACS, IHC-Fr, IP
Specificity	The clone 28-14-8 binds to the $\alpha$ 3 domain of H-2Db in the presence or absence of $\beta$ 2 microglobulin. It cross reacts with the $\alpha$ 3 domain of H-2Ld, but not Kd or Dd, and with H-2Dq and/or Lq. 28-14-8 can block H-2Ld-specific and H-2Ld-restricted antigen-specific lysis of target cells by cytotoxic T lymphocytes but it does not block recognition of H-2Ld-positive target cells by Ly-6G2-positive NK cells.
Host	Mouse
Clonality	Monoclonal
Clone	28-14-8
Isotype	IgG2a, kappa
Target Name	MHC Class I H2 Db
Species	Mouse
Immunogen	C3H.SW mouse splenocytes
Conjugation	Un-conjugated
Alternate Names	H-2 class I histocompatibility antigen, D-B alpha chain; H2-D; B; H-2D

## **Application Instructions**

Application table	Application	Dilution
	BL	Assay-dependent
	FACS	< 1 µg/10^6 cells
	IHC-Fr	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate reco should be determined by the	ommended starting dilutions and the optimal dilutions or concentrations e scientist.

## Properties

Form	Liquid
Purification Note	Low endotoxin
Buffer	PBS
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw

## Bioinformation

Note

Database links	GeneID: 14964 Mouse
	Swiss-port # P01899 Mouse
Gene Symbol	H2-D1
Gene Full Name	histocompatibility 2, D region locus 1
Function	Involved in the presentation of foreign antigens to the immune system. [UniProt]
Calculated Mw	41 kDa