

ARG21647 Goat anti-Human IgM antibody (AP), pre-adsorbed

Package: 500 µl
Store at: 4°C

Summary

Product Description	AP-conjugated Goat Polyclonal antibody recognizes Human IgM
Tested Reactivity	Hu
Tested Application	Depletion, ELISA, ELISPOT, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, Puri, WB
Specificity	The antibody reacts with the heavy chain of Human IgM. The antibody is pre-adsorbed with Human IgG and IgA, so the antibody may not react with Human IgG and IgA, but may react with IgM from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgM
Antigen Species	Human
Conjugation	AP

Application Instructions


Pre Adsorbed	Human IgG and IgA.																							
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>Depletion</td><td>Assay-dependent</td></tr><tr><td>ELISA</td><td>1:2000 - 1:4000</td></tr><tr><td>ELISPOT</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>Puri</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>		Application	Dilution	Depletion	Assay-dependent	ELISA	1:2000 - 1:4000	ELISPOT	Assay-dependent	FACS	Assay-dependent	FLISA	Assay-dependent	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	IHC-P	Assay-dependent	Puri	Assay-dependent	WB	Assay-dependent
Application	Dilution																							
Depletion	Assay-dependent																							
ELISA	1:2000 - 1:4000																							
ELISPOT	Assay-dependent																							
FACS	Assay-dependent																							
FLISA	Assay-dependent																							
ICC/IF	Assay-dependent																							
IHC-Fr	Assay-dependent																							
IHC-P	Assay-dependent																							
Puri	Assay-dependent																							
WB	Assay-dependent																							
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.																							

Properties

Form	Liquid
------	--------

Purification	Affinity purification with immunogen.
Buffer	50 mM Tris (pH 8.0), 0.1% Sodium azide, 1 mM MgCl ₂ and 50% Glycerol
Preservative	0.1% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21647 Goat anti-Human IgM antibody (AP) (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified Human IgM, IgG, and IgA. Immunoglobulins were detected with serially diluted ARG21647 Goat anti-Human IgM antibody (AP) (pre-adsorbed).