

**ARG21665**  
**Goat anti-Human IgD antibody, F(ab')<sub>2</sub> fragment, pre-adsorbed**Package: 500 µg  
Store at: -20°C

### Summary

Product Description	F(ab') <sub>2</sub> fragment of Goat Polyclonal antibody recognizes Human IgD
Tested Reactivity	Hu
Tested Application	ELISA, FACS, FLISA, IHC-Fr, IHC-P
Specificity	The antibody reacts with the heavy chain of Human IgD. The antibody is pre-adsorbed with Human IgG, IgM and IgA, so the antibody may not react with Human IgG, IgM and IgA, but may react with IgD from other species.
Host	Goat
Clonality	Polyclonal
Isotype	F(ab') <sub>2</sub> IgG
Target Name	IgD
Antigen Species	Human
Conjugation	Un-conjugated

### Application Instructions

Pre Adsorbed Human IgG, IgM and IgA.


**Application table**

Application	Dilution
ELISA	Assay-dependent
FACS	Assay-dependent
FLISA	Assay-dependent
IHC-Fr	Assay-dependent
IHC-P	Assay-dependent

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


### Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.



ARG21665 Goat anti-Human IgD antibody, F(ab')2 fragment (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified Human IgD, IgG, IgM, and IgA. Immunoglobulins were detected with serially diluted ARG21665 Goat anti-Human IgD antibody, F(ab')2 fragment (pre-adsorbed) followed by [ARG23857](#) Swine anti-Goat IgG (H+L) antibody (HRP) (pre-adsorbed).



ARG21665 Goat anti-Human IgD antibody, F(ab')2 fragment (pre-adsorbed) FACS image

Flow Cytometry: Human peripheral blood lymphocytes stained with ARG21665 Goat anti-Human IgD antibody, F(ab')2 fragment (pre-adsorbed) and [ARG21260](#) anti-CD19 antibody [SJ25-C1] (PE) followed by [ARG23856](#) Swine anti-Goat IgG (H+L) antibody (FITC) (pre-adsorbed).