

ARG21668 Goat anti-Human IgG antibody (TRITC), pre-adsorbed

Package: 500 µg
Store at: 4°C

Summary

Product Description	TRITC-conjugated Goat Polyclonal antibody recognizes Human IgG
Tested Reactivity	Hu
Tested Application	Depletion, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	The antibody reacts with the heavy chain of Human IgG. The antibody is pre-adsorbed with Human IgM and IgA, so the antibody may not react with Human IgM and IgA, but may react with IgG from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgG
Antigen Species	Human
Conjugation	TRITC

Application Instructions


Pre Adsorbed	Human IgM and IgA.																	
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>Depletion</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>1:100 - 1:400</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>		Application	Dilution	Depletion	Assay-dependent	FACS	Assay-dependent	FLISA	1:100 - 1:400	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	IHC-P	Assay-dependent	WB	Assay-dependent
Application	Dilution																	
Depletion	Assay-dependent																	
FACS	Assay-dependent																	
FLISA	1:100 - 1:400																	
ICC/IF	Assay-dependent																	
IHC-Fr	Assay-dependent																	
IHC-P	Assay-dependent																	
WB	Assay-dependent																	
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.																	

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide

Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21668 Goat anti-Human IgG antibody (TRITC) (pre-adsorbed)
FLISA image

FLISA: The plate was coated with purified Human IgG, IgM and IgA. Immunoglobulins were detected with serially diluted ARG21668 Goat anti-Human IgG antibody (TRITC) (pre-adsorbed).