

## ARG21719 Goat anti-Rabbit IgM antibody (TRITC), pre-adsorbed

Package: 500 µg  
Store at: 4°C

### Summary

|                     |   |
|---------------------|---|
| Product Description | TRITC-conjugated Goat Polyclonal antibody recognizes Rabbit IgM   |
| Tested Reactivity   | Rb  |
| Tested Application  | ELISA, ELISPOT, FACS, FLISA, IHC-Fr, IHC-P, WB  |
| Specificity         | The antibody reacts with the heavy chain of Rabbit IgM. The antibody is pre-adsorbed with Rabbit IgG, so the antibody may not react with Rabbit IgG, but may react with immunoglobulins from other species. |
| Host                | Goat  |
| Clonality           | Polyclonal  |
| Isotype             | IgG   |
| Target Name         | IgM   |
| Species             | Rabbit  |
| Conjugation         | TRITC   |

### Application Instructions

| Pre Adsorbed      | Rabbit IgG.  |                 |
|-------------------|--|-----------------|
| Application table | Application  | Dilution        |
|                   | ELISA  | Assay-dependent |
|                   | ELISPOT  | Assay-dependent |
|                   | FACS   | Assay-dependent |
|                   | FLISA  | 1:100 - 1:400   |
|                   | IHC-Fr   | Assay-dependent |
|                   | IHC-P  | Assay-dependent |
|                   | WB   | Assay-dependent |
| Application Note  | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |                 |

### Properties

|               |                                       |
|---------------|---------------------------------------|
| Form          | Liquid                                |
| Purification  | Affinity purification with immunogen. |
| Buffer        | PBS and 0.1% Sodium azide.            |
| Preservative  | 0.1% Sodium azide                     |
| Concentration | 1 mg/ml                               |

|                            |  |
|----------------------------|--|
| <b>Storage instruction</b> | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| <b>Note</b>                | For laboratory research only, not for drug, diagnostic or other use.   |