

ARG22008 anti-CD28 antibody [37.51] (FITC)

Package: 250 μg Store at: 4°C

Summary

Product Description	FITC-conjugated Syrian Hamster Monoclonal antibody [37.51] recognizes CD28
Tested Reactivity	Ms
Tested Application	BL, Cell-Act , FACS
Specificity	Mouse CD28
Host	Hamster
Clonality	Monoclonal
Clone	37.51
lsotype	lgG2
Target Name	CD28
Species	Mouse
Immunogen	C57BL/6N Mouse T cell lymphoma EL-4 cell line
Conjugation	FITC
Alternate Names	Tp44; TP44; T-cell-specific surface glycoprotein CD28; CD antigen CD28

Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Cell-Act	Assay-dependent
	FACS	< 2 µg/10^6 cells
Application Note	* The dilutions indicate recomm should be determined by the sci	nended starting dilutions and the optimal dilutions or concentrations ientist.


Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database linksGeneID: 12487 MouseSwiss-port # P31041 MouseGene SymbolCD28Gene Full NameCD28 antigenBackgroundThe protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]FunctionInvolved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. [UniProt]Research AreaDevelopmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibodyPTMCD40LG induces tyrosine phosphorylation of isoform 3.		
Gene SymbolCD28Gene Full NameCD28 antigenBackgroundThe protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]FunctionInvolved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. [UniProt]Research AreaDevelopmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibodyCalculated Mw25 kDa	Database links	GenelD: 12487 Mouse
Gene Full NameCD28 antigenBackgroundThe protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]FunctionInvolved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. [UniProt]Research AreaDevelopmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibodyCalculated Mw25 kDa		Swiss-port # P31041 Mouse
BackgroundThe protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]FunctionInvolved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. [UniProt]Research AreaDevelopmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibodyCalculated Mw25 kDa	Gene Symbol	CD28
and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]FunctionInvolved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. [UniProt]Research AreaDevelopmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibody 25 kDa	Gene Full Name	CD28 antigen
of T-cell survival. [UniProt] Research Area Developmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibody Calculated Mw 25 kDa	Background	and T-helper type-2 development. Several alternatively spliced transcript variants encoding different
Calculated Mw 25 kDa	Function	
	Research Area	Developmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibody
PTM CD40LG induces tyrosine phosphorylation of isoform 3.	Calculated Mw	25 kDa
	PTM	CD40LG induces tyrosine phosphorylation of isoform 3.

Images



ARG22008 anti-CD28 antibody [37.51] (FITC) FACS image

Flow Cytometry: BALB/c Mouse thymocytes stained with <u>ARG22008</u> anti-CD28 antibody [37.51] (FITC) and <u>ARG20819</u> anti-CD3e antibody [C363.29B] (PE).