

## ARG22101 anti-MHC Class II I Ek antibody [17-3-3S] (FITC)

Package: 250 µg  
Store at: 4°C

### Summary

Product Description	FITC-conjugated Mouse Monoclonal antibody [17-3-3S] recognizes MHC Class II I Ek
Tested Reactivity	Ms
Tested Application	FACS, IHC-Fr
Specificity	Mouse I-Ek
Host	Mouse
Clonality	Monoclonal
Clone	17-3-3S
Isotype	IgG2a, kappa
Target Name	MHC Class II I Ek
Species	Mouse
Immunogen	C3H Mouse skin graft and splenocytes
Conjugation	FITC
Alternate Names	AI323765; H-2Ea; MHC-H2-Ea; H2-Ea; I-Ealpha; H-2 class II histocompatibility antigen, E-U alpha chain; Ia3; E-alpha-f; Ia-3

### Application Instructions


Application table	Application	Dilution
	FACS	< 1 µg/10 <sup>6</sup> cells
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Database links	<a href="#">GeneID: 100504404 Mouse</a> <a href="#">Swiss-port # P14439 Mouse</a>
Gene Symbol	H2-Ea-ps
Gene Full Name	histocompatibility 2, class II antigen E alpha, pseudogene
Background	This locus belongs to the class II major histocompatibility complex (MHC) family of genes, which encode immune response (Ia) antigens that function in the T-cell-dependent immune response. This family member has multiple haplotypes, some of which result in the production of an E-alpha subunit that combines with an E-beta subunit to form a functional E complex at the cell surface. Other haplotypes, including that of the reference genome allele, contain mutations and they thus represent polymorphic pseudogenes that do not produce functional products. These mutations include frameshifting indels, nonsense mutations, and deletions of larger regions. The reference genome haplotype contains a deletion at the 5' end of the gene, including the core promoter region and the transcription start site, and therefore no transcripts result from this haplotype. [provided by RefSeq, Aug 2011]
Calculated Mw	29 kDa

Images



ARG22101 anti-MHC Class II I Ek antibody [17-3-3S] (FITC) FACS image

Flow Cytometry: AKR Mouse splenocytes stained with [ARG22101](#) anti-MHC Class II I Ek antibody [17-3-3S] (FITC) and [ARG20819](#) anti-CD3e antibody [C363.29B] (PE).