

Product datasheet

info@arigobio.com

ARG22156 anti-CD3 epsilon antibody [UCHT1] (PE-Cyanine 7)

Package: 50 tests Store at: 4°C

Summary

Product Description PE-Cyanine 7-conjugated Mouse Monoclonal antibody [UCHT1] recognizes CD3 epsilon

Tested Reactivity Hu

Tested Application BL, Cell-Act , FACS, ICC/IF, IHC-Fr, IHC-P, WB

Specificity Human CD3 epsilon

Host Mouse

Clonality Monoclonal

Clone UCHT1

Target Name IgG1, kappa
CD3 epsilon

Species Human

Immunogen Human peripheral blood lymphocytes from a patient with Sézary's syndrome and infant thymocytes

Conjugation PE-Cyanine 7

Alternate Names CD3E; CD3 Epsilon Subunit Of T-Cell Receptor Complex; T-Cell Surface Glycoprotein CD3 Epsilon Chain;

CD3e Antigen, Epsilon Polypeptide (TiT3 Complex); T-Cell Surface Antigen T3/Leu-4 Epsilon Chain; CD3e Molecule, Epsilon (CD3-TCR Complex); CD3-Epsilon; CD3epsilon; T3E; T-Cell Antigen Receptor Complex,

Epsilon Subunit Of T3; CD3e Molecule; CD3e Antigen; CD3-EPSILON; CD3EPSILON

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|------------------|
| | BL | Assay-dependent |
| | Cell-Act | Assay-dependent |
| | FACS | 10 μl/10^6 cells |
| | ICC/IF | Assay-dependent |
| | IHC-Fr | Assay-dependent |
| | IHC-P | Assay-dependent |
| | WB | Assay-dependent |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

| Form | Liquid |
|--------|-------------------------------------|
| Buffer | PBS, 0.1% Sodium azide and Sucrose. |

Preservative 0.1% Sodium azide

Stabilizer Sucrose

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links <u>GeneID: 915 Human</u>

Swiss-port # P04234 Human

Gene Symbol CD3E

Gene Full Name CD3 Epsilon Subunit Of T-Cell Receptor Complex

Background The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma,

-delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also

been linked to a susceptibility to type I diabetes in women.

Function Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in

adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their

cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein

tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathway.

Highlight Related products:

CD3 antibodies; CD3 ELISA Kits; CD3 Duos / Panels;

Related news:

New antibody panels and duos for Tumor immune microenvironment

Tumor-Infiltrating Lymphocytes (TILs)

Research Area Cancer antibody; Developmental Biology antibody; Immune System antibody; Lymphocyte Marker


antibody; Inflammatory Cell Marker antibody; T-cell Marker antibody; T-cell infiltration Study antibody;

Tumor-infiltrating Lymphocyte Study antibody

Calculated Mw 23 kDa

PTM Disulfide bond, Phosphoprotein

Cellular Localization Cell membrane, Membrane



ARG22156 anti-CD3 epsilon antibody [UCHT1] (PE-Cyanine 7) FACS image

Flow Cytometry: Human peripheral blood lymphocytes stained with ARG22156 anti-CD3 epsilon antibody [UCHT1] (PE-Cyanine 7) and ARG21260 anti-CD19 antibody [SJ25-C1] (PE).