

## ARG22256 anti-TCP1 alpha antibody [91a]

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rat Monoclonal antibody [91a] recognizes TCP1 alpha
Tested Reactivity	Hu, Ms, Rat, Bov, Ce, Dm, Dog, Gpig, Hm, Mk, Pig, Rb, Yeast
Tested Application	FACS, ICC/IF, IP, WB
Specificity	Detects ~60kDa. Also detects ~92kDa. Cross reactivity with human HSP60 has been observed with this antibody in immunoblot analysis. Reacts weakly with <i>Saccharomyces cerevisiae</i> .
Host	Rat
Clonality	Monoclonal
Clone	91a
Isotype	IgG2a
Target Name	TCP1 alpha
Species	Mouse
Immunogen	Recombinant Mouse TCP1 alpha protein fragment (carboxy terminal region).
Conjugation	Un-conjugated
Alternate Names	T-complex protein 1 subunit alpha; CCT1; D6S230E; TCP-1-alpha; CCTa; CCT-alpha

### Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	ICC/IF	1:100
	IP	Assay-dependent
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~60 kDa	

### Properties


Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS (pH 7.4), 0.1% Sodium azide and 50% Glycerol
Preservative	0.1% Sodium azide
Stabilizer	50% Glycerol

Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	Tcp1
Gene Full Name	t-complex protein 1
Background	The protein encoded by this gene is a molecular chaperone that is a member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin. Alternate transcriptional splice variants of this gene, encoding different isoforms, have been characterized. In addition, three pseudogenes that appear to be derived from this gene have been found. [provided by RefSeq, Jun 2010]
Function	Molecular chaperone; assists the folding of proteins upon ATP hydrolysis. As part of the BBS/CCT complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia. Known to play a role, in vitro, in the folding of actin and tubulin. [UniProt]
Calculated Mw	60 kDa
Cellular Localization	Cytoplasm

## Images




ARG22256 anti-TCP1 alpha antibody [91a] ICC/IF image

Immunocytochemistry: 2% Formaldehyde (20 min at RT) fixed Heat Shocked HeLa cells stained with ARG22256 anti-TCP1 alpha antibody [91a] (yellow) at 1:100 dilution (12 hours at 4°C). Counterstain: DAPI (blue) nuclear stain at 1:40000 for 120 min at RT. Magnification: 100x. Left: DAPI (blue) nuclear stain, Middle: Primary antibody, Right: Composite.



ARG22256 anti-TCP1 alpha antibody [91a] WB image

Western blot: 1) MW ladder, 20 µg of 2) A431 lysate, and 3) HEK293 lysate stained with ARG22256 anti-TCP1 alpha antibody [91a] (60 min at RT). Block: 5% milk + TBST 1hr at RT.



ARG22256 anti-TCP1 alpha antibody [91a] ICC/IF image

Immunocytochemistry: 2% Formaldehyde (20 min at RT) fixed Heat Shocked HeLa cells stained with ARG22256 anti-TCP1 alpha antibody [91a] (red) at 1:100 dilution (12 hours at 4°C). Counterstain: DAPI (blue) nuclear stain at 1:40000 for 120 min at RT. Magnification: 20x. Left: DAPI (blue) nuclear stain, Middle: Primary antibody, Right: Composite.

---