

ARG23181 anti-CD16.2 / FcγRIV antibody [AT137]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Rat Monoclonal antibody [AT137] recognizes CD16.2 / FcγRIV Mouse anti human CD16-2 antibody, clone AT137, recognizes CD16-2, also known as FcRIV. CD16-2 is a low affinity activatory receptor that binds to IgG2a and IgG2b with intermediate affinity. FcRIV has no affinity for IgG1 or IgG3 and therefore has a unique subclass specificity when compared to other Fc receptors. Expression of CD16-2 is restricted to cells of the myeloid lineage.
Tested Reactivity	Ms
Tested Application	FACS, IHC-Fr
Host	Rat
Clonality	Monoclonal
Clone	AT137
Isotype	IgG2a
Target Name	CD16.2 / FcγRIV
Species	Mouse
Immunogen	Fusion Protein consisting of mouse CD16-2 and rat CD4.
Conjugation	Un-conjugated
Alternate Names	Fcrl3; CD16-2; FcγRIV; Fcγr3a; FcγammaRIV; 4833442P21Rik

Application Instructions

Application table	Application	Dilution
	FACS	1:10
	IHC-Fr	1:100 - 1:1000
Application Note	FACS: Use 10 µl of the suggested working dilution to label 10 ⁶ cells in 100 µl. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	TRIS buffered saline and 0.09% Sodium azide.
Preservative	0.09% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	Fcgr4
Gene Full Name	Fc receptor, IgG, low affinity IV
Function	Receptor for the Fc region of immunoglobulin gamma (PubMed:16039578). Also acts as a receptor for the Fc region of immunoglobulin epsilon (PubMed:17558411, PubMed:18949059). Binds with intermediate affinity to both IgG2a and IgG2b (PubMed:16039578, PubMed:17558411, PubMed:19795417). Can bind to IgG2a and IgG2b monomers (PubMed:18949059). Does not display binding to IgG1 or IgG3 (PubMed:16039578). Mediates neutrophil activation by IgG complexes redundantly with Fcgr3 (PubMed:18097064). Plays a role in promoting bone resorption by enhancing osteoclast differentiation following binding to IgG2a (PubMed:25824719). Binds with low affinity to both the a and b allotypes of IgE (PubMed:18949059). Has also been shown to bind to IgE allotype a only but not to allotype b (PubMed:17558411). Binds aggregated IgE but not the monomeric form and bound monomeric IgG is readily displaced by IgE complexes (PubMed:18949059). Binding to IgE promotes macrophage-mediated phagocytosis, antigen presentation to T cells, production of proinflammatory cytokines and the late phase of cutaneous allergic reactions (PubMed:17558411, PubMed:18949059). [UniProt]
Calculated Mw	28 kDa