

ARG23211 anti-hCG (Holo C3 epitope) antibody [INN-hCG-45]

Package: 250 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [INN-hCG-45] recognizes hCG (Holo C3 epitope) Mouse anti hCG (Holo C3 Epitope) antibody, clone INN-hCG-45 recognizes intact human chorionic gonadotropin (hCG), binding to the c3 epitope on hCG. Mouse anti hCG (Holo C3 Epitope) antibody, clone INN-hCG-45 does not recognize the free α or β subunits of human chorionic gonadotropin.
Tested Reactivity	Hu
Tested Application	ELISA, RIA
Host	Mouse
Clonality	Monoclonal
Clone	INN-hCG-45
Isotype	IgG1
Target Name	hCG (Holo C3 epitope)
Species	Human
Immunogen	Human CG.
Conjugation	Un-conjugated
Alternate Names	hCGB; CGB5; CGB7; CGB3; Chorionic gonadotrophin chain beta; CGB8; CG-beta; Choriogonadotropin subunit beta

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	RIA	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS and 0.09% Sodium azide.
Preservative	0.09% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CGB
Gene Full Name	chorionic gonadotropin, beta polypeptide
Background	This gene is a member of the glycoprotein hormone beta chain family and encodes the beta 3 subunit of chorionic gonadotropin (CG). Glycoprotein hormones are heterodimers consisting of a common alpha subunit and an unique beta subunit which confers biological specificity. CG is produced by the trophoblastic cells of the placenta and stimulates the ovaries to synthesize the steroids that are essential for the maintenance of pregnancy. The beta subunit of CG is encoded by 6 genes which are arranged in tandem and inverted pairs on chromosome 19q13.3 and contiguous with the luteinizing hormone beta subunit gene. [provided by RefSeq, Jul 2008]
Function	Stimulates the ovaries to synthesize the steroids that are essential for the maintenance of pregnancy. [UniProt]
Calculated Mw	18 kDa