

Summary

ARG23227 anti-TNF alpha antibody [MP9-20A4] (PE)

Package: 50 tests Store at: 4°C

Product Description	PE-conjugated Rat Monoclonal antibody [MP9-20A4] recognizes TNF alpha Rat anti Human TNF alpha antibody, clone MP9-20A4 recognizes human tumor Necrosis Factor - alpha (TNF alpha), also known as TNFSF2. This ~17kDa cytokine is expressed by activated macrophages, monocytes, neutrophils, T cells and NK cells. Rat anti Human TNF alpha antibody, clone MP9-20A4 is reported to act as a blocking antibody (Abrams et al. 2001).
Tested Reactivity	Hu
Tested Application	FACS
Host	Rat
Clonality	Monoclonal
Clone	MP9-20A4
Isotype	lgG1
Target Name	TNF alpha
Species	Human
Conjugation	PE
Alternate Names	Tumor necrosis factor ligand superfamily member 2; DIF; Cachectin; ICD2; ICD1; N-terminal fragment; TNF-a; TNFA; TNFSF2; TNF-alpha; Tumor necrosis factor; NTF

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
Application Note	FACS: Membrane permeabilisation is required for this application. Use 10 μl of the suggested working dilution to label 10^6 cells in 100 μl. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified
Buffer	PBS, 0.09% Sodium azide, 1% BSA and 5% Sucrose.
Preservative	0.09% Sodium azide
Stabilizer	1% BSA and 5% Sucrose
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol	TNF
Gene Full Name	tumor necrosis factor
Background	This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine. [provided by RefSeq, Jul 2008]
Function	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation. Impairs regulatory T-cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Upregulates the expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective. Key mediator of cell death in the anticancer action of BCG-stimulated neutrophils in combination with DIABLO/SMAC mimetic in the RT4v6 bladder cancer cell line.
	The TNF intracellular domain (ICD) form induces IL12 production in dendritic cells. [UniProt]
Highlight	Related products: <u>TNF alpha antibodies; TNF alpha ELISA Kits; TNF alpha Duos / Panels; TNF alpha recombinant</u> <u>proteins; Anti-Rat IgG secondary antibodies;</u> Related news: <u>HMGB1 in inflammation</u> <u>Inflammatory Cytokines</u>
Calculated Mw	26 kDa
ΡΤΜ	The soluble form derives from the membrane form by proteolytic processing. The membrane-bound form is further proteolytically processed by SPPL2A or SPPL2B through regulated intramembrane proteolysis producing TNF intracellular domains (ICD1 and ICD2) released in the cytosol and TNF C-domain 1 and C-domain 2 secreted into the extracellular space.
	The membrane form, but not the soluble form, is phosphorylated on serine residues. Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1.
	O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid. [UniProt]