

ARG23357
anti-IL2 antibody [B-G5] (FITC)Package: 500 µl
Store at: 4°C

Summary

Product Description	FITC-conjugated Mouse Monoclonal antibody [B-G5] recognizes IL2
Tested Reactivity	Hu
Tested Application	FACS
Specificity	This antibody recognizes both natural and recombinant human IL-2.
Host	Mouse
Clonality	Monoclonal
Clone	B-G5
Isotype	IgG1
Target Name	IL2
Species	Human
Immunogen	Natural human IL-2
Conjugation	FITC
Alternate Names	TCGF; IL-2; lymphokine; Interleukin-2; Aldesleukin; T-cell growth factor

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
Application Note	FACS: Use 10 µl of antibody to label 5 x 10 ⁵ cells. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Buffer	PBS, 0.1% Sodium azide and 5% BSA.
Preservative	0.1% Sodium azide
Stabilizer	5% BSA
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	IL2
Gene Full Name	interleukin 2
Background	The protein encoded by this gene is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli. [provided by RefSeq, Jul 2008]
Function	Produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells. [UniProt]
Calculated Mw	18 kDa