

ARG24001 anti-Collagen I antibody

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Collagen I
Tested Reactivity	Hu, Bov
Tested Application	ELISA, ICC/IF, IHC-P
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Collagen I
Species	Bovine
Immunogen	Type I collagen extracted from Bovine skin.
Conjugation	Un-conjugated
Alternate Names	OI1; OI2; OI3; OI4; EDSC; Collagen alpha-1(I) chain; Alpha-1 type I collagen

Application Instructions

Application table	Application	Dilution
	ELISA	1:2000
	ICC/IF	1:40
	IHC-P	1:500
Application Note	IHC-P: Antigen Retrieval: treat sections with 0.5% hyaluronidase at RT for 1hr before staining. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Purified.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	COL1A1
Gene Full Name	collagen, type I, alpha 1

Background	<p>This gene encodes the pro-alpha1 chains of type I collagen whose triple helix comprises two alpha1 chains and one alpha2 chain. Type I is a fibril-forming collagen found in most connective tissues and is abundant in bone, cornea, dermis and tendon. Mutations in this gene are associated with osteogenesis imperfecta types I-IV, Ehlers-Danlos syndrome type VIIA, Ehlers-Danlos syndrome Classical type, Caffey Disease and idiopathic osteoporosis. Reciprocal translocations between chromosomes 17 and 22, where this gene and the gene for platelet-derived growth factor beta are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans, resulting from unregulated expression of the growth factor. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene. [provided by R. Dalglish, Feb 2008]</p>
Function	Type I collagen is a member of group I collagen (fibrillar forming collagen). [UniProt]
Highlight	<p>Related products: Collagen I antibodies; Collagen I ELISA Kits; Collagen I Duos / Panels; Anti-Rabbit IgG secondary antibodies;</p> <p>Related news: New antibody panels for Myofibroblasts and CAFs</p>
Calculated Mw	139 kDa
PTM	<p>Proline residues at the third position of the tripeptide repeating unit (G-X-P) are hydroxylated in some or all of the chains. Proline residues at the second position of the tripeptide repeating unit (G-P-X) are hydroxylated in some of the chains.</p> <p>O-linked glycan consists of a Glc-Gal disaccharide bound to the oxygen atom of a post-translationally added hydroxyl group. [UniProt]</p>
Cellular Localization	Secreted, extracellular space, extracellular matrix. [UniProt]

Images



ARG24001 anti-Collagen I antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Bovine skin tissue stained with ARG24001 anti-Collagen I antibody.