

Product datasheet

info@arigobio.com

ARG30162 Cytotoxic T Cell Surface Marker Antibody Panel (FACS) Package: 1 kit Store at: -20°C, 4°C

Component

Cat. No.	Component Name	Host clonality	Reactivity	Application	Package
ARG62928	anti-CD8 antibody [MEM-31]	Mouse mAb	Hu	CyTOF®-candidate, FACS, IP	50 μg
ARG62889	anti-CD54 / ICAM1 antibody [1H4] (FITC)	Mouse mAb	Hu	FACS	50 tests
ARG53814	anti-CD28 antibody [CD28.2] (APC)	Mouse mAb	Hu, NHuPrm	FACS	50 tests
ARG62855	anti-CD45 antibody [MEM-28] (Biotin)	Mouse mAb	Hu	FACS	50 μg

Summary

Product Description A cytotoxic T cell is a T lymphocyte that kills cancer cells, virally infected cells and cells that are under

damage. Most T lymphocytes express a subset of surface markers such as CD8, CD45 and CD54. CD28 expresses on the surface of T cells and provide co-stimulatory signals required for T cell activation.

Target Name Cytotoxic T Cell Surface Marker

Alternate Names Cytotoxic T Cell Surface Marker antibody; APC-conjugated CD28 antibody; Biotin-conjugated CD45

antibody; FITC-conjugated CD54 / ICAM1 antibody; CD8 antibody

Properties

Storage instruction Store antibodies at 4°C or -20°C. Please refer to the each product datasheet for detail temperatures of

the antibodies.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Full Name Antibody Panel for Cytotoxic T Cell Surface Marker


Highlight Related Product: anti-CD8 antibody;

anti-CD54 / ICAM1 antibody;

anti-CD28 antibody; anti-CD45 antibody;

Research Area Cell Biology and Cellular Response antibody; Developmental Biology antibody; Immune System


antibody; Neuroscience antibody; Signaling Transduction antibody



Monoclonal antibody clone 1H4 Flow Cytometry analysis image


Flow Cytometry: U937 human histiocytic lymphoma cell stained with antibody clone 1H4.

Total viable cells were used for analysis.




Monoclonal antibody clone CD28.2 Flow Cytometry analysis image

Flow Cytometry: Human peripheral blood leukocytes stained with antibody clone CD28.2.




Monoclonal antibody clone MEM-28 Flow Cytometry analysis image

Flow Cytometry: Human peripheral blood cells stained with antibody clone MEM-28.




Monoclonal antibody clone MEM-28 ICC/IF image

Immunofluorescence: Human peripheral blood mononuclear cell stained with clone MEM-28 (green)
Cell nuclei was stained with DAPI (blue).




Monoclonal antibody clone MEM-31 Flow Cytometry analysis image

Flow Cytometry: Human peripheral blood cells stained with antibody clone MEM-31.




ARG62928 anti-CD8 antibody [MEM-31] CyTOF image

CyTOF: PBMC (after FicoII-Paque separation) stained with ARG62928 anti-CD8 antibody [MEM-31] (Sm152). Singlet cells were gated for data analysis.




ARG62889 anti-CD54 / ICAM1 antibody [1H4] (FITC) FACS image

Flow Cytometry: Separation of Human CD54 positive Monocytes (red) from Human CD54 negative Lymphocytes (black-dashed). Human peripheral blood stained with ARG62889 anti-CD54 / ICAM1 antibody [1H4] (FITC).



ARG53814 anti-CD28 antibody [CD28.2] (APC) FACS image

Flow Cytometry: Human peripheral blood leukocytes stained with ARG53814 anti-CD28 antibody [CD28.2] (APC).



ARG62855 anti-CD45 antibody [MEM-28] (Biotin) FACS image

Flow Cytometry: Human peripheral blood cells stained with ARG62855 anti-CD45 antibody [MEM-28] (Biotin), followed by Streptavidin (APC).