

## ARG40040 anti-PSMA6 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes PSMA6
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PSMA6
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-246 of Human PSMA6 (NP_002782.1).
Conjugation	Un-conjugated
Alternate Names	Proteasome subunit alpha type-6; Proteasome iota chain; PROS27; EC 3.4.25.1; PROS-27; 27 kDa prosomal protein; p27K; IOTA; Macropain iota chain; Multicatalytic endopeptidase complex iota chain

### Application Instructions


Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse skeletal muscle and MCF7	
Observed Size	30 kDa	

### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Gene Symbol	PSMA6
Gene Full Name	proteasome subunit alpha 6
Background	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene. A pseudogene has been identified on the Y chromosome. [provided by RefSeq, Aug 2013]
Function	The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. [UniProt]
Calculated Mw	27 kDa
Cellular Localization	Cytoplasm. Nucleus. Note=Colocalizes with TRIM5 in cytoplasmic bodies. [UniProt]

Images



ARG40040 anti-PSMA6 antibody WB image

Western blot: 25 µg of Mouse skeletal muscle and MCF7 cell lysates stained with ARG40040 anti-PSMA6 antibody at 1:1000 dilution.



ARG40040 anti-PSMA6 antibody WB image

Western blot: 25 µg of Mouse brain, PC-12 and A549 cell lysates stained with ARG40040 anti-PSMA6 antibody at 1:1000 dilution through one-step method.