

## ARG40108 anti-PCM1 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes PCM1
Tested Reactivity	Hu, Ms
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PCM1
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-260 of Human PCM1 (NP_006188.3).
Conjugation	Un-conjugated
Alternate Names	Pericentriolar material 1 protein; RET/PCM-1; hPCM-1; PTC4; PCM-1

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Raji	
Observed Size	280 kDa	

### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	PCM1
Gene Full Name	pericentriolar material 1
Background	The protein encoded by this gene is a component of centriolar satellites, which are electron dense granules scattered around centrosomes. Inhibition studies show that this protein is essential for the correct localization of several centrosomal proteins, and for anchoring microtubules to the centrosome. Chromosomal aberrations involving this gene are associated with papillary thyroid carcinomas and a variety of hematological malignancies, including atypical chronic myeloid leukemia and T-cell lymphoma. [provided by RefSeq, Sep 2010]
Function	Required for centrosome assembly and function. Essential for the correct localization of several centrosomal proteins including CEP250, CETN3, PCNT and NEK2. Required to anchor microtubules to the centrosome. Involved in the biogenesis of cilia. [UniProt]
Calculated Mw	229 kDa
PTM	Ubiquitinated. Undergoes monoubiquitination catalyzed by the E3 ubiquitin-protein ligase MIB1 in proliferating cells, preventing cilia formation. Monoubiquitination by MIB1 is inhibited in response to cellular stress, such as ultraviolet light (UV) radiation or heat shock, resulting in cilia formation initiation. [UniProt]
Cellular Localization	Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasmic granule. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite. Cytoplasm, cytoskeleton, cilium basal body. Note=Recruitment to the centrosome requires microtubules and dynein. [UniProt]

Images



ARG40108 anti-PCM1 antibody WB image

Western blot: 25 µg of Raji cell lysate stained with ARG40108 anti-PCM1 antibody at 1:1000 dilution.