

# Product datasheet

info@arigobio.com

# ARG40502 anti-NTR3 / Sortilin antibody

Package: 100 μl Store at: -20°C

### Summary

Product Description Rabbit Polyclonal antibody recognizes NTR3 / Sortilin

Tested Reactivity Hu, Ms, Rat
Tested Application IHC-P, WB
Host Rabbit
Clonality Polyclonal

Isotype IgG

Target Name NTR3 / Sortilin

Species Human

Immunogen Synthetic peptide derived from Human NTR3 / Sortilin.

Conjugation Un-conjugated

Alternate Names Neurotensin receptor 3; Sortilin; 100 kDa NT receptor; NT3; NTR3; Gp95; LDLCQ6; Glycoprotein 95

#### **Application Instructions**

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

#### **Properties**

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol SORT1

Gene Full Name sortilin 1

Background This gene encodes a protein that is a multi-ligand type-1 receptor with similarity to the yeast

carboxypeptidase Y sorting receptor Vps10 protein. The encoded protein, a trans-Golgi network (TGN) transmembrane protein, binds a number of unrelated ligands that participate in a wide range of cellular processes; however, it lacks the typical features of a signalling receptor. In the TGN, furin mediates the activation of the mature binding form. The encoded protein consists of a large luminal domain, a single transmembrane segment and short C-terminal cytoplasmic tail. The luminal domain contains a cysteinerich region similar to two corresponding segments in the yeast Vps10p; the cytoplasmic tail is similar to the corresponding segment of the cation-independent mannose 6-phosphate receptor and the tail also interacts with the VHS domains of GGA (Golgi-associated, gamma-adaptin homologous, ARF-

interacting) proteins. [provided by RefSeq, Jul 2008]

Function Functions as a sorting receptor in the Golgi compartment and as a clearance receptor on the cell

surface. Required for protein transport from the Golgi apparatus to the lysosomes by a pathway that is independent of the mannose-6-phosphate receptor (M6PR). Also required for protein transport from the Golgi apparatus to the endosomes. Promotes neuronal apoptosis by mediating endocytosis of the proapoptotic precursor forms of BDNF (proBDNF) and NGFB (proNGFB). Also acts as a receptor for neurotensin. May promote mineralization of the extracellular matrix during osteogenic differentiation by scavenging extracellular LPL. Probably required in adipocytes for the formation of specialized storage vesicles containing the glucose transporter SLC2A4/GLUT4 (GLUT4 storage vesicles, or GSVs). These vesicles provide a stable pool of SLC2A4 and confer increased responsiveness to insulin. May also

mediate transport from the endoplasmic reticulum to the Golgi. [UniProt]

Calculated Mw 92 kDa


PTM The N-terminal propeptide is cleaved by furin and possibly other homologous proteases. [UniProt]

Cellular Localization Membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus, Golgi membrane protein. Endosome membrane; Single-pass type I membrane protein.

stack membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Cell membrane; Single-pass type I membrane protein; Extracellular side. Lysosome

membrane; Single-pass type I membrane protein. [UniProt]

#### **Images**



## ARG40502 anti-NTR3 / Sortilin antibody WB image

Western blot: SW480 cell lysate stained with ARG40502 anti-NTR3 / Sortilin antibody.