

# Product datasheet

info@arigobio.com

ARG41059 anti-USF1 antibody

Package: 100 μl Store at: -20°C

### **Summary**

Isotype

**Product Description** Rabbit Polyclonal antibody recognizes USF1

IgG

**Tested Reactivity** Hu, Ms, Rat **Tested Application** IHC-P, WB Host Rabbit Polyclonal

Clonality

USF1 **Target Name** 

**Species** Human

Immunogen Recombinant fusion protein corresponding to aa. 1-310 of Human USF1 (NP\_009053.1).

Conjugation Un-conjugated

**Alternate Names** HYPLIP1; FCHL1; MLTFI; bHLHb11; UEF; Upstream stimulatory factor 1; Class B basic helix-loop-helix

protein 11; Major late transcription factor 1; FCHL; MLTF

## **Application Instructions**

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse kidney	
Observed Size	41 kDa	

## **Properties**

Form Liquid

Purification Affinity purified.

**Buffer** PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot Storage instruction

> and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

For laboratory research only, not for drug, diagnostic or other use. Note

#### Bioinformation

Gene Symbol USF1

Gene Full Name upstream transcription factor 1

Background This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a

cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL). Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has

been defined on chromosome 21. [provided by RefSeq, Feb 2013]


Function Transcription factor that binds to a symmetrical DNA sequence (E-boxes) (5'-CACGTG-3') that is found in

a variety of viral and cellular promoters. [UniProt]

Calculated Mw 34 kDa


Cellular Localization Nucleus. [UniProt]

## **Images**



#### ARG41059 anti-USF1 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human kidney cancer stained with ARG41059 anti-USF1 antibody at 1:200 dilution.



## ARG41059 anti-USF1 antibody WB image

Western blot: 25  $\mu g$  of Mouse kidney lysate stained with ARG41059 anti-USF1 antibody at 1:1000 dilution.