

ARG41085 anti-Coilin antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Coilin
Tested Reactivity	Hu
Tested Application	FACS, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Coilin
Species	Human
Immunogen	Synthetic peptide derived from Human Coilin.
Conjugation	Un-conjugated
Alternate Names	Coilin; p80-coilin; CLN80

Application Instructions

Application table	Application	Dilution
	FACS	1:100
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	77 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	COIL
Gene Full Name	coilin
Background	The protein encoded by this gene is an integral component of Cajal bodies (also called coiled bodies). Cajal bodies are nuclear suborganelles of varying number and composition that are involved in the post-transcriptional modification of small nuclear and small nucleolar RNAs. The N-terminus of the coilin protein directs its self-oligomerization while the C-terminus influences the number of nuclear bodies assembled per cell. Differential methylation and phosphorylation of coilin likely influences its localization among nuclear bodies and the composition and assembly of Cajal bodies. This gene has pseudogenes on chromosome 4 and chromosome 14. [provided by RefSeq, Jul 2008]
Function	Component of nuclear coiled bodies, also known as Cajal bodies or CBs, which are involved in the modification and assembly of nucleoplasmic snRNPs. [UniProt]
Calculated Mw	63 kDa
PTM	Symmetrical dimethylation of arginine residues within the RG repeat region enhances affinity for SMN, and thus localization of SMN complexes to CBs. Phosphorylation during mitosis is associated with disassembly of CBs. [UniProt]
Cellular Localization	Nucleus. Nucleus, Cajal body. [UniProt]

Images



ARG41085 anti-Coilin antibody WB image

Western blot: Jurkat cell lysate stained with ARG41085 anti-Coilin antibody.