

## Product datasheet

info@arigobio.com

# ARG41457 anti-PTH / Parathyroid Hormone antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Product Description Rabbit Polyclonal antibody recognizes PTH / Parathyroid Hormone

Tested Reactivity Hu

Tested Application IHC-P, WB

Host Rabbit

**Clonality** Polyclonal

Isotype IgG

Target Name PTH / Parathyroid Hormone

Species Human

Immunogen Synthetic peptide of Human PTH / Parathyroid Hormone.

Conjugation Un-conjugated

Alternate Names PTH1; Parathyroid hormone; Parathyrin; Parathormone; PTH

### **Application Instructions**

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Human PTH Recombinant protein	
Observed Size	~ 37 kDa	

### **Properties**

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol PTH

Gene Full Name parathyroid hormone

Background The protein encoded by this gene is a hormone secreted by parathyroid cells. This hormone elevates

blood Ca2+ level by dissolving the salts in bone and preventing their renal excretion. Defects in this gene are a cause of familial isolated hypoparathyroidism (FIH). [provided by RefSeq, Jul 2008]

Function PTH elevates calcium level by dissolving the salts in bone and preventing their renal excretion.

Stimulates [1-14C]-2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblastic cells.

[UniProt]

Calculated Mw 13 kDa

Cellular Localization Secreted. [UniProt]