

ARG41610 anti-RPA2 / RPA32 antibody [RPPA-1]

Package: 50 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [RPPA-1] recognizes RPA2 / RPA32
Tested Reactivity	Hu
Tested Application	WB
Host	Mouse
Clonality	Monoclonal
Clone	RPPA-1
Isotype	IgG2b, kappa
Target Name	RPA2 / RPA32
Species	Human
Immunogen	Full length recombinant Human RPA2 / RPA32 protein.
Conjugation	Un-conjugated
Alternate Names	RF-A protein 2; RPA32; RP-A p34; Replication protein A 34 kDa subunit; Replication factor A protein 2; Replication protein A 32 kDa subunit; REPA2; RP-A p32; RFA2

Application Instructions

Application table	Application	Dilution
	WB	1 - 2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 30 kDa	


Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS, 0.05% Sodium azide and 0.1 mg/ml BSA.
Preservative	0.05% Sodium azide
Stabilizer	0.1 mg/ml BSA
Concentration	0.2 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol	RPA2
Gene Full Name	replication protein A2, 32kDa
Function	As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response. It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair. Plays also a role in base excision repair (BER) probably through interaction with UNG. Through RFW3 may activate CHEK1 and play a role in replication checkpoint control. Also recruits SMARCA1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance. [UniProt]
Calculated Mw	29 kDa
PTM	Differentially phosphorylated throughout the cell cycle, becoming phosphorylated at the G1-S transition and dephosphorylated in late mitosis. Mainly phosphorylated at Ser-23 and Ser-29, by cyclin A-CDK2 and cyclin B-CDK1, respectively during DNA replication and mitosis. Dephosphorylation may require the serine/threonine-protein phosphatase 4. Phosphorylation at Ser-23 and Ser-29 is a prerequisite for further phosphorylation. Becomes hyperphosphorylated on additional residues including Ser-4, Ser-8, Thr-21 and Ser-33 in response to DNA damage. Hyperphosphorylation is mediated by ATM, ATR and PRKDC. Primarily recruited to DNA repair nuclear foci as a hypophosphorylated form it undergoes subsequent hyperphosphorylation, catalyzed by ATR. Hyperphosphorylation is required for RAD51 recruitment to chromatin and efficient DNA repair. Phosphorylation at Thr-21 depends upon RFW3 presence. DNA damage-induced 'Lys-63'-linked polyubiquitination by PRPF19 mediates ATRIP recruitment to the RPA complex at sites of DNA damage and activation of ATR. [UniProt]
Cellular Localization	Nucleus. Nucleus, PML body. Note=Redistributes to discrete nuclear foci upon DNA damage in an ATR-dependent manner. [UniProt]

Images



ARG41610 anti-RPA2 / RPA32 antibody [RPPA-1] WB image

Western blot: A375 cell lysate stained with ARG41610 anti-RPA2 / RPA32 antibody [RPPA-1].