

## ARG41864 anti-FUBP1 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes FUBP1
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	FUBP1
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 302-644 of Human FUBP1. (NP_003893.2)
Conjugation	Un-conjugated
Alternate Names	FBP; Far upstream element-binding protein 1; FUSE-binding protein 1; hDH V; FUBP; DNA helicase V

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	IP	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	U-87MG	
Observed Size	79 kDa	


### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation


Gene Symbol	FUBP1
Gene Full Name	far upstream element (FUSE) binding protein 1
Background	<p>The protein encoded by this gene is a single stranded DNA-binding protein that binds to multiple DNA elements, including the far upstream element (FUSE) located upstream of c-myc. Binding to FUSE occurs on the non-coding strand, and is important to the regulation of c-myc in undifferentiated cells. This protein contains three domains, an amphipathic helix N-terminal domain, a DNA-binding central domain, and a C-terminal transactivation domain that contains three tyrosine-rich motifs. The N-terminal domain is thought to repress the activity of the C-terminal domain. This protein is also thought to bind RNA, and contains 3'-5' helicase activity with in vitro activity on both DNA-DNA and RNA-RNA duplexes. Aberrant expression of this gene has been found in malignant tissues, and this gene is important to neural system and lung development. Binding of this protein to viral RNA is thought to play a role in several viral diseases, including hepatitis C and hand, foot and mouth disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014]</p>
Function	<p>Regulates MYC expression by binding to a single-stranded far-upstream element (FUSE) upstream of the MYC promoter. May act both as activator and repressor of transcription. [UniProt]</p>
Calculated Mw	68 kDa
PTM	Ubiquitinated. This targets the protein for proteasome-mediated degradation. [UniProt]
Cellular Localization	Nucleus. [UniProt]

## Images




ARG41864 anti-FUBP1 antibody ICC/IF image

Immunofluorescence: C6 cells stained with ARG41864 anti-FUBP1 antibody at 1:100 dilution.




ARG41864 anti-FUBP1 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human lung cancer tissue stained with ARG41864 anti-FUBP1 antibody 1:100 dilution.



#### ARG41864 anti-FUBP1 antibody WB image

Western blot: 25 µg of U-87MG cell lysate stained with ARG41864 anti-FUBP1 antibody at 1:1000 dilution.



#### ARG41864 anti-FUBP1 antibody IP image

Immunoprecipitation: 200 µg extracts of HeLa cells were immunoprecipitated and stained with ARG41864 anti-FUBP1 antibody at 1:1000 dilution.