

ARG42065 anti-VEGF antibody (Biotin)

Package: 25 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Goat Polyclonal antibody recognizes VEGF
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	VEGF
Species	Human
Immunogen	Recombinant Human VEGF.
Conjugation	Biotin
Alternate Names	MVCD1; Vascular permeability factor; VEGF-A; VPF; VEGF; Vascular endothelial growth factor A

Application Instructions

Application table	Application	Dilution
	ELISA	0.25 - 1 µg/ml
	WB	0.1 - 0.2 µg/ml
Application Note	Sandwich ELISA (Capture antibody - Detection antibody): ARG42064 (0.5 - 2 µg/ml) - ARG42065 (0.25 - 1 µg/ml) * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	VEGFA
Gene Full Name	vascular endothelial growth factor A
Function	Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth. Binding to NRP1 receptor initiates a signaling pathway needed for motor neuron axon guidance and cell body migration, including for the caudal migration of facial motor neurons from rhombomere 4 to rhombomere 6 during embryonic development (By similarity). [UniProt]
Highlight	<p>Related products:</p> <p>VEGF antibodies; VEGF ELISA Kits; VEGF Duos / Panels; VEGF recombinant proteins; Anti-Goat IgG secondary antibodies;</p> <p>Related news:</p> <p>The role of HDGF in tumor angiogenesis</p>
Calculated Mw	22 kDa
Cellular Localization	Secreted. Note=VEGF121 is acidic and freely secreted. VEGF165 is more basic, has heparin-binding properties and, although a significant proportion remains cell-associated, most is freely secreted. VEGF189 is very basic, it is cell-associated after secretion and is bound avidly by heparin and the extracellular matrix, although it may be released as a soluble form by heparin, heparinase or plasmin. [UniProt]

Images



ARG42065 anti-VEGF antibody (Biotin) ELISA image

Sandwich ELISA: Detecting human VEGF (using 100 µl/well antibody solution). [ARG42064](#) anti-VEGF antibody as a capture antibody at 0.5 - 2 µg/ml dilution in combination with ARG42065 anti-VEGF antibody (Biotin) as a detection antibody at 0.25 - 1.0 µg/ml dilution. Results of a typical standard run with optical density reading at 405 - 650 nm.