

ARG42069 anti-PRKRA antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes PRKRA
Tested Reactivity	Hu
Predict Reactivity	Bov, Rat
Tested Application	FACS, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PRKRA
Species	Human
Immunogen	KLH-conjugated synthetic peptide between aa. 225-252 of Human PRKRA.
Conjugation	Un-conjugated
Alternate Names	PACT; DYT16; Protein kinase, interferon-inducible double-stranded RNA-dependent activator; HSD14; Interferon-inducible double-stranded RNA-dependent protein kinase activator A; PKR-associating protein X; Protein activator of the interferon-induced protein kinase; PKR-associated protein X; RAX

Application Instructions

Application table	Application	Dilution
	FACS	1:10 - 1:50
	IHC-P	1:50 - 1:100
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HL-60	
Observed Size	~ 34 kDa	

Properties

Form	Liquid
Purification	Purification with Protein A and immunogen peptide.
Buffer	PBS and 0.09% (W/V) Sodium azide.
Preservative	0.09% (W/V) Sodium azide.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	PRKRA
Gene Full Name	protein kinase, interferon-inducible double stranded RNA dependent activator
Background	This gene encodes a protein kinase activated by double-stranded RNA which mediates the effects of interferon in response to viral infection. Mutations in this gene have been associated with dystonia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2008]
Function	Activates EIF2AK2/PKR in the absence of double-stranded RNA (dsRNA), leading to phosphorylation of EIF2S1/EIF2-alpha and inhibition of translation and induction of apoptosis. Required for siRNA production by DICER1 and for subsequent siRNA-mediated post-transcriptional gene silencing. Does not seem to be required for processing of pre-miRNA to miRNA by DICER1. Promotes UBC9-p53/TP53 association and sumoylation and phosphorylation of p53/TP53 at 'Lys-386' at 'Ser-392' respectively and enhances its activity in a EIF2AK2/PKR-dependent manner (By similarity). [UniProt]
Calculated Mw	34 kDa
PTM	Phosphorylated at Ser-246 in unstressed cells and at Ser-287 in stressed cells. Phosphorylation at Ser-246 appears to be a prerequisite for subsequent phosphorylation at Ser-287. Phosphorylation at Ser-246 and Ser-287 are necessary for activation of EIF2AK2/PKR under conditions of stress. [UniProt]
Cellular Localization	Cytoplasm, perinuclear region. Cytoplasm. [UniProt]

Images




ARG42069 anti-PRKRA antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human placenta tissue stained with ARG42069 anti-PRKRA antibody.



ARG42069 anti-PRKRA antibody WB image

Western blot: 35 µg of HL-60 cell lysate stained with ARG42069 anti-PRKRA antibody.



ARG42069 anti-PRKRA antibody FACS image

Flow Cytometry: HL-60 cells stained with ARG42069 anti-PRKRA antibody (bottom histogram) or without primary antibody as control (top histogram), followed by incubation with FITC labelled secondary antibody.