

ARG42473 anti-Adipophilin antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Adipophilin
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Adipophilin
Species	Human
Immunogen	Recombinant protein corresponding to K226-Q418 of Human Adipophilin.
Conjugation	Un-conjugated
Alternate Names	ADRP; Adipophilin; Adipose differentiation-related protein; ADFP; Perilipin-2

Application Instructions

Application table	Application	Dilution
	IHC-P	1:200 - 1:1000
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	MCF7	
Observed Size	~ 48 kDa	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na ₂ HPO ₄ , 0.9% NaCl, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol	PLIN2
Gene Full Name	perilipin 2
Background	The protein encoded by this gene belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This protein is associated with the lipid globule surface membrane material, and maybe involved in development and maintenance of adipose tissue. However, it is not restricted to adipocytes as previously thought, but is found in a wide range of cultured cell lines, including fibroblasts, endothelial and epithelial cells, and tissues, such as lactating mammary gland, adrenal cortex, Sertoli and Leydig cells, and hepatocytes in alcoholic liver cirrhosis, suggesting that it may serve as a marker of lipid accumulation in diverse cell types and diseases. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Mar 2011]
Function	May be involved in development and maintenance of adipose tissue. [UniProt]
Calculated Mw	48 kDa
PTM	Acylated; primarily with C14, C16 and C18 fatty acids. [UniProt]
Cellular Localization	Membrane; Peripheral membrane protein. Lipid droplet. [UniProt]


Images



ARG42473 anti-Adipophilin antibody IHC-P image

Immunohistochemistry: Human epithelial cells stained with ARG42473 anti-Adipophilin antibody.

From Satoshi Sumida et al. The Journal of Medical Investigation (2023), [doi: 10.2152/jmi.70.251](https://doi.org/10.2152/jmi.70.251), Fig. 2M.




ARG42473 anti-Adipophilin antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Rat testis tissue stained with ARG42473 anti-Adipophilin antibody at 1 μg/ml dilution.



ARG42473 anti-Adipophilin antibody WB image

Western blot: MCF7 whole cell lysate stained with ARG42473 anti-Adipophilin antibody at 0.5 µg/ml dilution.



ARG42473 anti-Adipophilin antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse testis tissue stained with ARG42473 anti-Adipophilin antibody at 1 µg/ml dilution.