

Product datasheet

info@arigobio.com

ARG42781 anti-Catalase antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes Catalase

Tested Reactivity Hu, Ms, Rat

Tested Application FACS, ICC/IF, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name Catalase

Species Mouse

Immunogen Recombinant protein of Mouse Catalase.

Conjugation Un-conjugated

Alternate Names Catalase; EC 1.11.1.6

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|----------|
| | FACS | 1:20 |
| | ICC/IF | 1:20 |
| | WB | 1:1000 |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |
| Positive Control | K562 | |
| Observed Size | ~ 62 kDa | |

Properties

Form Liquid

Purification Affinity purified.

Buffer 50 mM Tris-Glycine (pH 7.4), 150 mM NaCl, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.

Preservative 0.01% Sodium azide

Stabilizer 40% Glycerol and 0.05% BSA

Concentration Batch dependent

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

www.arigobio.com arigo, nuts about antibodies 1/2

Bioinformation

Gene Symbol CAT

Gene Full Name catalase

Background This gene encodes catalase, a key antioxidant enzyme in the bodies defense against oxidative stress.

Catalase is a heme enzyme that is present in the peroxisome of nearly all aerobic cells. Catalase converts the reactive oxygen species hydrogen peroxide to water and oxygen and thereby mitigates the toxic effects of hydrogen peroxide. Oxidative stress is hypothesized to play a role in the development of many chronic or late-onset diseases such as diabetes, asthma, Alzheimer's disease, systemic lupus erythematosus, rheumatoid arthritis, and cancers. Polymorphisms in this gene have been associated with decreases in catalase activity but, to date, acatalasemia is the only disease known to be caused by

this gene. [provided by RefSeq, Oct 2009]

Function Occurs in almost all aerobically respiring organisms and serves to protect cells from the toxic effects of


hydrogen peroxide. Promotes growth of cells including T-cells, B-cells, myeloid leukemia cells, melanoma cells, mastocytoma cells and normal and transformed fibroblast cells. [UniProt]

Calculated Mw 60 kDa

PTM The N-terminus is blocked. [UniProt]

Cellular Localization Peroxisome. [UniProt]

Images



ARG42781 anti-Catalase antibody WB image

Western blot: K562 cell lysate stained with ARG42781 anti-Catalase antibody at 1:1000 dilution.