

Product datasheet

info@arigobio.com

ARG43196 anti-VIPR1 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes VIPR1

Tested Reactivity Hu, Ms, Rat

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name VIPR1

Species Human

Immunogen Synthetic peptide derived from Human VIPR1.

Conjugation Un-conjugated

Alternate Names PACAP-R2; Pituitary adenylate cyclase-activating polypeptide type II receptor; V1RG; Vasoactive

intestinal polypeptide receptor 1; VIPR; VPAC1; VIP-R-1; RDC1; VAPC1; VPAC1R; II; PACAP type II

receptor; VPCAP1R; VIRG; HVR1; PACAP-R-2

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A549, H9C2 and Raw264.7	
Observed Size	~ 60 kDa	

Properties

Form Liquid

Purification Affinity purification with immunogen.

Purity > 95% (by SDS-PAGE)

Buffer PBS (pH 7.2), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

www.arigobio.com argo.nuts about antibodies 1/2

Bioinformation

Gene Symbol VIPR1

Gene Full Name vasoactive intestinal peptide receptor 1

Background This gene encodes a receptor for vasoactive intestinal peptide, a small neuropeptide. Vasoactive

intestinal peptide is involved in smooth muscle relaxation, exocrine and endocrine secretion, and water and ion flux in lung and intestinal epithelia. Its actions are effected through integral membrane receptors associated with a guanine nucleotide binding protein which activates adenylate cyclase. Several transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Oct 2011]


Function This is a receptor for VIP. The activity of this receptor is mediated by G proteins which activate adenylyl

cyclase. The affinity is VIP = PACAP-27 > PACAP-38. [UniProt]

Calculated Mw 52 kDa

Cellular Localization Cell membrane; Multi-pass membrane protein. [UniProt]

Images



ARG43196 anti-VIPR1 antibody WB image

Western blot: A549, H9C2 and Raw264.7 cell lysates stained with ARG43196 anti-VIPR1 antibody at 1:500 dilution.