

ARG43647 anti-MMP2 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes MMP2
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MMP2
Species	Human
Immunogen	Recombinant protein fragment corresponding to c-terminal region of Human MMP2.
Conjugation	Un-conjugated
Alternate Names	CLG4A; MMP-2; TBE-1; MONA; CLG4; EC 3.4.24.24; Gelatinase A; Matrix metalloproteinase-2; MMP-II; 72 kDa gelatinase; 72 kDa type IV collagenase

Application Instructions


Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	72 kDa	

Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS, 0.01% Thimerosal and 50% Glycerol.
Preservative	0.01% Thimerosal
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Gene Symbol	MMP2
Gene Full Name	matrix metalloproteinase 2
Background	<p>This gene is a member of the matrix metalloproteinase (MMP) gene family, that are zinc-dependent enzymes capable of cleaving components of the extracellular matrix and molecules involved in signal transduction. The protein encoded by this gene is a gelatinase A, type IV collagenase, that contains three fibronectin type II repeats in its catalytic site that allow binding of denatured type IV and V collagen and elastin. Unlike most MMP family members, activation of this protein can occur on the cell membrane. This enzyme can be activated extracellularly by proteases, or, intracellularly by its S-glutathiolation with no requirement for proteolytical removal of the pro-domain. This protein is thought to be involved in multiple pathways including roles in the nervous system, endometrial menstrual breakdown, regulation of vascularization, and metastasis. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2014]</p>
Function	<p>Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly- -Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. Involved in the formation of the fibrovascular tissues in association with MMP14.</p> <p>PEX, the C-terminal non-catalytic fragment of MMP2, possesses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrinα5β3 on the surface of blood vessels.</p> <p>Isoform 2: Mediates the proteolysis of CHUK/IκBα and initiates a primary innate immune response by inducing mitochondrial-nuclear stress signaling with activation of the pro-inflammatory NF-κB, NFAT and IRF transcriptional pathways. [UniProt]</p>
Calculated Mw	74 kDa
PTM	<p>Phosphorylation on multiple sites modulates enzymatic activity. Phosphorylated by PKC in vitro. The propeptide is processed by MMP14 (MT-MMP1) and MMP16 (MT-MMP3). Autocatalytic cleavage in the C-terminal produces the anti-angiogenic peptide, PEX. This processing appears to be facilitated by binding integrinα5β3.</p>
Cellular Localization	Cytoplasm; Extracellular matrix; Membrane; Mitochondrion; Nucleus; Secreted

Images




ARG43647 anti-MMP2 antibody WB image

Western blot: U-87 MG stained with ARG43647 anti-MMP2 antibody at 1:1000 dilution.




ARG43647 anti-MMP2 antibody ICC/IF image

Immunofluorescence: PC-12 stained with ARG43647 anti-MMP2 antibody at 1:100 dilution.




ARG43647 anti-MMP2 antibody WB image

Western blot: Rat lung stained with ARG43647 anti-MMP2 antibody at 1:1000 dilution.



ARG43647 anti-MMP2 antibody IHC-P image

Immunohistochemistry: Mouse lung stained with ARG43647 anti-MMP2 antibody at 1:100 dilution.



ARG43647 anti-MMP2 antibody ICC/IF image

Immunofluorescence: NIH/3T3 stained with ARG43647 anti-MMP2 antibody at 1:100 dilution.

ARG43647 anti-MMP2 antibody WB image

Western blot: C2C12 stained with ARG43647 anti-MMP2 antibody at 1:1000 dilution.

